



Vaasan yliopisto
UNIVERSITY OF VAASA

D. BAIDIUK

Completion and extension of operators in Krein spaces

PROCEEDINGS OF THE UNIVERSITY OF VAASA
WORKING PAPERS 10
MATHEMATICS 6

Publisher Vaasan yliopisto		Date of publication February 2016	
Author D. Baidiuk	Type of publication Working papers		
	Name and number of series Proceedings of the University of Vaasa. Working Papers, 10 Proceedings of the University of Vaasa. Working papers, Mathematics, 6		
Contact information University of Vaasa Faculty of Technology Department of Mathematics and Statistics P.O. Box 700 FI-65101 Vaasa Finland	ISBN 978-952-476-668-5 (online)		
	ISSN 1799-7658 (Proceedings of the University of Vaasa. Working Papers 10, online)		
	Number of pages 23		Language English
Title of publication Completion and extension of operators in Kreĭn spaces			
Abstract A generalization of the well-known results of M.G. Krein about the description of selfadjoint contractive extension of a hermitian contraction is obtained. This generalization concerns the situation, where the selfadjoint operator A and extensions \widetilde{A} belong to a Krein space or a Pontryagin space and their defect operators are allowed to have a fixed number of negative eigenvalues. Also a result of Yu.L. Shmul'yan on completions of nonnegative block operators is generalized for block operators with a fixed number of negative eigenvalues in a Krein space. This paper is a natural continuation of S. Hassi's and author's paper [5].			
Keywords Completion, extension of operators, Krein and Pontryagin spaces MSC 2010 Primary 46C20, 47A20, 47A63; Secondary 47B25			

Contents

1. Introduction	1
2. A completion problem for block operators in Kreĭn spaces	3
3. Some inertia formulas	5
4. A pair of completion problems in a Kreĭn space	7
5. Completion problem in a Pontryagin space	10
5.1. Defect operators and link operators	10
5.2. Lemmas on negative indices of certain block operators	11
5.3. Contractive extensions of contractions with minimal negative indices	15
References	19

1. Introduction

In 1947 M.G. Kreĭn published one of his famous papers [17] on a description of a nonnegative selfadjoint extensions of a densely defined nonnegative operator A in a Hilbert space. Namely, all nonnegative selfadjoint extensions \tilde{A} of A can be characterized by the following two inequalities:

$$(A_F + a)^{-1} \leq (\tilde{A} + a)^{-1} \leq (A_K + a)^{-1}, \quad a > 0,$$

where the Friedrichs (hard) extension A_F and the Kreĭn-von Neumann (soft) extension A_K of A . He proved these results by transforming the problems the study of contractive operators.

The first result of the present paper is a generalization of a result due to Shmul'yan [19] on completions of nonnegative block operators where the result was applied for introducing so-called Hellinger operator integrals. This result was extended in [5] for block operators in a Hilbert space by allowing a fixed number of negative eigenvalues. In Section 2 this result is further extended to block operators which act in a Kreĭn space.

In paper [5] we studied classes of "quasi-contractive" symmetric operators T_1 allowing a finite number of negative eigenvalues for the associated defect operator $I - T_1^* T_1$, i.e., $\nu_-(I - T_1^* T_1) < \infty$ as well as "quasi-nonnegative" operators A with $\nu_-(A) < \infty$ and the existence and description of all possible selfadjoint extensions T and \tilde{A} of them which preserve the given negative indices $\nu_-(I - T^2) = \nu_-(I - T_1^* T_1)$ and $\nu_-(\tilde{A}) = \nu_-(A)$, and proved precise analogs of the above mentioned results of M.G. Kreĭn under a minimality condition on the negative indices $\nu_-(I - T_1^* T_1)$ and $\nu_-(A)$, respectively. It was an unexpected fact that when there is a solution then the solution set still contains a minimal solution and a maximal solution which then describe the whole solution set via two operator inequalities, just as in the original paper of M.G. Kreĭn. In this paper analogous results are established for "quasi-contractive" operators acting in a Kreĭn space; see Theorems 4.2, 5.7.

In Section 4 a first Kreĭn space analog of completion problem is formulated and a description of its solutions is found. Namely, we consider classes of "quasi-contractive" symmetric operators T_1 in a Kreĭn space with $\nu_-(I - T_1^* T_1) < \infty$ and we describe all possible selfadjoint (in the Kreĭn space sense) extensions T of T_1 which preserve the given negative index $\nu_-(I - T^* T) = \nu_-(I - T_1^* T_1)$. This problem is close to the completion problem studied in [5] and has a similar description for its solutions. For further history behind this problem see also [1, 2, 3, 7, 8, 9, 10, 11, 12, 14, 15, 16, 20].

The main result of the present paper is proved in Section 5. Namely, we consider classes of "quasi-contractive" symmetric operators T_1 in a Kreĭn space (\mathfrak{H}, J) with

$$\nu_-[I - T_1^{[*]} T_1] := \nu_-(J(I - T_1^{[*]} T_1)) < \infty \quad (1.1)$$

and we establish a solvability criterion and a description of all possible selfadjoint extensions T of T_1 (in the Kreĭn space sense) which preserve the given negative index $\nu_-[I - T^{[*]} T] = \nu_-[I - T_1^{[*]} T_1]$. It should be pointed out that

in this more general setting the descriptions involve so-called link operator L_T which was introduced by Arsene, Constantintscu and Gheondea in [3] (see also [2, 7, 8, 18]).

2. A completion problem for block operators in Kreĭn spaces

By definition the modulus $|C|$ of a closed operator C is the nonnegative selfadjoint operator $|C| = (C^*C)^{1/2}$. Every closed operator admits a polar decomposition $C = U|C|$, where U is a (unique) partial isometry with the initial space $\overline{\text{ran}}|C|$ and the final space $\overline{\text{ran}}C$, cf. [13]. For a selfadjoint operator $H = \int_{\mathbb{R}} t dE_t$ in a Hilbert space \mathfrak{H} the partial isometry U can be identified with the signature operator, which can be taken to be unitary: $J = \text{sign}(H) = \int_{\mathbb{R}} \text{sign}(t) dE_t$, in which case one should define $\text{sign}(t) = 1$ if $t \geq 0$ and otherwise $\text{sign}(t) = -1$.

Let \mathcal{H} be a Hilbert space, and let $J_{\mathcal{H}}$ be a signature operator in it, i.e., $J_{\mathcal{H}} = J_{\mathcal{H}}^* = J_{\mathcal{H}}^{-1}$. We interpret the space \mathcal{H} as a Kreĭn space $(\mathcal{H}, J_{\mathcal{H}})$ (see [4, 6]) in which the indefinite scalar product is defined by the equality

$$[\varphi, \psi]_{\mathcal{H}} = (J_{\mathcal{H}}\varphi, \psi)_{\mathcal{H}}.$$

Let us introduce a partial ordering for selfadjoint Kreĭn space operators. For selfadjoint operators A and B with the same domains $A \geq_J B$ if and only if $[(A - B)f, f] \geq 0$ for all $f \in \text{dom } A$. If not otherwise indicated the word "smallest" means the smallest operator in the sense of this partial ordering.

Consider a bounded incomplete block operator

$$A^0 = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & * \end{pmatrix} \begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix} \rightarrow \begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix} \quad (2.1)$$

in the Kreĭn space $\mathfrak{H} = (\mathfrak{H}_1 \oplus \mathfrak{H}_2, J)$, where (\mathfrak{H}_1, J_1) and (\mathfrak{H}_2, J_2) are Kreĭn spaces with fundamental symmetries J_1 and J_2 , and $J = \begin{pmatrix} J_1 & 0 \\ 0 & J_2 \end{pmatrix}$.

Theorem 2.1. *Let $\mathfrak{H} = (\mathfrak{H}_1 \oplus \mathfrak{H}_2, J)$ be an orthogonal decomposition of the Kreĭn space \mathfrak{H} and let A^0 be an incomplete block operator of the form (2.1). Assume that $A_{11} = A_{11}^{[*]}$ and $A_{21} = A_{12}^{[*]}$ are bounded, the numbers of negative squares of the quadratic form $[A_{11}f, f]$ ($f \in \text{dom } A_{11}$) $\nu_-[A_{11}] := \nu_-(J_1 A_{11}) = \kappa < \infty$, where $\kappa \in \mathbb{Z}_+$, and let us introduce $J_{11} := \text{sign}(J_1 A_{11})$ the (unitary) signature operator of $J_1 A_{11}$. Then:*

- (i) *There exists a completion $A \in [(\mathfrak{H}, J)]$ of A^0 with some operator $A_{22} = A_{22}^{[*]} \in [(\mathfrak{H}_2, J_2)]$ such that $\nu_-[A] = \nu_-[A_{11}] = \kappa$ if and only if*

$$\text{ran } J_1 A_{12} \subset \text{ran } |A_{11}|^{1/2}.$$

- (ii) *In this case the operator $S = |A_{11}|^{[-1/2]} J_1 A_{12}$, where $|A_{11}|^{[-1/2]}$ denotes the (generalized) Moore-Penrose inverse of $|A_{11}|^{1/2}$, is well defined and $S \in [(\mathfrak{H}_2, J_2), (\mathfrak{H}_1, J_1)]$. Moreover, $S^{[*]} J_1 J_{11} S$ is the "smallest" operator in the solution set*

$$\mathcal{A} := \left\{ A_{22} = A_{22}^{[*]} \in [(\mathfrak{H}_2, J_2)] : A = (A_{ij})_{i,j=1}^2 : \nu_-[A] = \kappa \right\}$$

and this solution set admits a description

$$\mathcal{A} = \left\{ A_{22} \in [(\mathfrak{H}_2, J_2)] : A_{22} = J_2(S^* J_{11} S + Y) = S^{[*]} J_1 J_{11} S + J_2 Y, \right. \\ \left. \text{where } Y = Y^* \geq 0 \right\}.$$

Proof. Let us introduce a block operator

$$\tilde{A}^0 = \begin{pmatrix} \tilde{A}_{11} & \tilde{A}_{12} \\ \tilde{A}_{21} & * \end{pmatrix} = \begin{pmatrix} J_1 A_{11} & J_1 A_{12} \\ J_2 A_{21} & * \end{pmatrix}.$$

The blocks of this operator satisfy the identities $\tilde{A}_{11} = \tilde{A}_{11}^*$, $\tilde{A}_{21}^* = \tilde{A}_{12}$ and

$$\begin{aligned} \text{ran } J_1 A_{11} &= \text{ran } \tilde{A}_{11} \subset \text{ran } |\tilde{A}_{11}|^{1/2} = \text{ran } (\tilde{A}_{11}^* \tilde{A}_{11})^{1/4} \\ &= \text{ran } (A_{11}^* A_{11})^{1/4} = \text{ran } |A_{11}|^{1/2}. \end{aligned}$$

Then due to [5, Theorem 1] a description of all selfadjoint operator completions of \tilde{A}^0 admits representation $\tilde{A} = \begin{pmatrix} \tilde{A}_{11} & \tilde{A}_{12} \\ \tilde{A}_{21} & \tilde{A}_{22} \end{pmatrix}$ with $\tilde{A}_{22} = \tilde{S}^* J_{11} \tilde{S} + Y$, where $\tilde{S} = |\tilde{A}_{11}|^{[-1/2]} \tilde{A}_{12}$ and $Y = Y^* \geq 0$.

This yields description for the solutions of the completion problem. The set of completions has the form $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$, where

$$\begin{aligned} A_{22} &= J_2 \tilde{A}_{22} = J_2 A_{21} J_1 |A_{11}|^{[-1/2]} J_{11} |A_{11}|^{[-1/2]} J_1 A_{12} + J_2 Y \\ &= J_2 S^* J_{11} S + J_2 Y = S^{[*]} J_1 J_{11} S + J_2 Y. \end{aligned}$$

□

3. Some inertia formulas

Some simple inertia formulas are now recalled. The factorization $H = B^{[*]}EB$ clearly implies that $\nu_{\pm}[H] \leq \nu_{\pm}[E]$, cf. (1.1). If H_1 and H_2 are selfadjoint operators in a Kreĭn space, then

$$H_1 + H_2 = \begin{pmatrix} I \\ I \end{pmatrix}^{[*]} \begin{pmatrix} H_1 & 0 \\ 0 & H_2 \end{pmatrix} \begin{pmatrix} I \\ I \end{pmatrix}$$

shows that $\nu_{\pm}[H_1 + H_2] \leq \nu_{\pm}[H_1] + \nu_{\pm}[H_2]$. Consider the selfadjoint block operator $H \in [(\mathfrak{H}_1, J_1) \oplus (\mathfrak{H}_2, J_2)]$, where $J_i = J_i^* = J_i^{-1}$, ($i = 1, 2$) of the form

$$H = H^{[*]} = \begin{pmatrix} A & B^{[*]} \\ B & I \end{pmatrix},$$

By applying the above mentioned inequalities shows that

$$\nu_{\pm}[A] \leq \nu_{\pm}[A - B^{[*]}B] + \nu_{\pm}(J_2). \quad (3.1)$$

Assuming that $\nu_{-}[A - B^{[*]}J_2B]$ and $\nu_{-}(J_2)$ are finite, the question when $\nu_{-}[A]$ attains its maximum in (3.1), or equivalently, $\nu_{-}[A - B^{[*]}J_2B] \geq \nu_{-}[A] - \nu_{-}(J_2)$ attains its minimum, turns out to be of particular interest. The next result characterizes this situation as an application of Theorem 2.1. Recall that if $J_1A = J_A|A|$ is the polar decomposition of J_1A , then one can interpret $\mathfrak{H}_A = (\overline{\text{ran}} J_1A, J_A)$ as a Kreĭn space generated on $\overline{\text{ran}} J_1A$ by the fundamental symmetry $J_A = \text{sign}(J_1A)$.

Theorem 3.1. *Let $A \in [(\mathfrak{H}_1, J_1)]$ be selfadjoint, $B \in [(\mathfrak{H}_1, J_1), (\mathfrak{H}_2, J_2)]$, $J_i = J_i^* = J_i^{-1} \in [\mathfrak{H}_i]$, ($i = 1, 2$), and assume that $\nu_{-}[A], \nu_{-}(J_2) < \infty$. If the equality*

$$\nu_{-}[A] = \nu_{-}[A - B^{[*]}B] + \nu_{-}(J_2)$$

holds, then $\text{ran } J_1B^{[]} \subset \text{ran } |A|^{1/2}$ and $J_1B^{[*]} = |A|^{1/2}K$ for a unique operator $K \in [(\mathfrak{H}_2, J_2), \mathfrak{H}_A]$ which is J -contractive: $J_2 - K^*J_AK \geq 0$.*

Conversely, if $B^{[]} = |A|^{1/2}K$ for some J -contractive operator $K \in [(\mathfrak{H}_2, J_2), \mathfrak{H}_A]$, then the equality (3.1) is satisfied.*

Proof. Assume that (3.1) is satisfied. The factorization

$$H = \begin{pmatrix} A & B^{[*]} \\ B & I \end{pmatrix} = \begin{pmatrix} I & B^{[*]} \\ 0 & I \end{pmatrix} \begin{pmatrix} A - B^{[*]}B & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} I & 0 \\ B & I \end{pmatrix}$$

shows that $\nu_{-}[H] = \nu_{-}[A - B^{[*]}B] + \nu_{-}(J_2)$, which combined with the equality (3.1) gives $\nu_{-}[H] = \nu_{-}[A]$. Therefore, by Theorem 2.1 one has $\text{ran } J_1B^{[*]} \subset \text{ran } |A|^{1/2}$ and this is equivalent to the existence of a unique operator $K \in [(\mathfrak{H}_2, J_2), \mathfrak{H}_A]$ such that $J_1B^{[*]} = |A|^{1/2}K$; i.e. $K = |A|^{[-1/2]}J_1B^{[*]}$. Furthermore, $K^{[*]}J_1J_AK \leq_{J_2} I$ by the minimality property of $K^{[*]}J_1J_AK$ in Theorem 2.1, in other words K is a J -contraction.

Converse, if $J_1B^{[*]} = |A|^{1/2}K$ for some J -contractive operator $K \in [(\mathfrak{H}_2, J_2), \mathfrak{H}_A]$, then clearly $\text{ran } J_1B^{[*]} \subset \text{ran } |A|^{1/2}$. By Theorem 2.1 the completion problem for H^0 has solutions with the minimal solution $S^{[*]}J_1J_AS$,

where

$$S = |A|^{[-1/2]} J_1 B^{[*]} = |A|^{[-1/2]} |A|^{1/2} K = K.$$

Furthermore, by J -contractivity of K one has $K^{[*]} J_1 J_A K \leq_{J_2} I$, i.e. I is also a solution and thus $\nu_-[H] = \nu_-[A]$ or, equivalently, the equality (3.1) is satisfied. \square

4. A pair of completion problems in a Kreĭn space

In this section we introduce and describe the solutions of a Kreĭn space version of a completion problem that was treated in [5].

Let $(\mathfrak{H}_i, (J_i \cdot, \cdot))$ and $(\mathfrak{H}, (J \cdot, \cdot))$ be Kreĭn spaces, where $\mathfrak{H} = \mathfrak{H}_1 \oplus \mathfrak{H}_2$, $J = \begin{pmatrix} J_1 & 0 \\ 0 & J_2 \end{pmatrix}$, and J_i are fundamental symmetries ($i = 1, 2$), let $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ be an operator such that $\nu_-(I - T_{11}^* T_{11}) = \kappa < \infty$. Denote $\tilde{T}_{11} = J_1 T_{11}$, then $\tilde{T}_{11} = \tilde{T}_{11}^*$ in the Hilbert space \mathfrak{H}_1 . Rewrite $\nu_-(I - T_{11}^* T_{11}) = \nu_-(I - \tilde{T}_{11}^2)$. Denote

$$J_+ = \text{sign}(I - \tilde{T}_{11}), \quad J_- = \text{sign}(I + \tilde{T}_{11}), \quad \text{and} \quad J_{11} = \text{sign}(I - \tilde{T}_{11}^2),$$

and let $\kappa_+ = \nu_-(J_+)$ and $\kappa_- = \nu_-(J_-)$. It is easy to get that $J_{11} = J_- J_+ = J_+ J_-$. Moreover, there is an equality $\kappa = \kappa_- + \kappa_+$ (see [5, Lemma 5.1]). We recall the results for the operator \tilde{T}_{11} from the paper [5] and after that reformulate them for the operator T_{11} . We recall completion problem and its solutions that was investigated in a Hilbert space setting in [5]. The problem concerns the existence and a description of selfadjoint operators \tilde{T} such that $\tilde{A}_+ = I + \tilde{T}$ and $\tilde{A}_- = I - \tilde{T}$ solve the corresponding completion problems

$$\tilde{A}_\pm^0 = \begin{pmatrix} I \pm \tilde{T}_{11} & \pm \tilde{T}_{21}^* \\ \pm \tilde{T}_{21} & * \end{pmatrix}, \quad (4.1)$$

under *minimal index conditions* $\nu_-(I + \tilde{T}) = \nu_-(I + \tilde{T}_{11})$, $\nu_-(I - \tilde{T}) = \nu_-(I - \tilde{T}_{11})$, respectively. The solution set is denoted by $\text{Ext}_{\tilde{T}_{11}, \kappa}(-1, 1)$.

The next theorem gives a general solvability criterion for the completion problem (4.1) and describes all solutions to this problem.

Theorem 4.1. ([5, Theorem 5]) *Let $\tilde{T}_1 = \begin{pmatrix} \tilde{T}_{11} \\ \tilde{T}_{21} \end{pmatrix} : \mathfrak{H}_1 \rightarrow \begin{pmatrix} \mathfrak{H}_1 \\ \mathfrak{H}_2 \end{pmatrix}$ be a symmetric operator with $\tilde{T}_{11} = \tilde{T}_{11}^* \in [\mathfrak{H}_1]$ and $\nu_-(I - \tilde{T}_{11}^2) = \kappa < \infty$, and let $J_{11} = \text{sign}(I - \tilde{T}_{11}^2)$. Then the completion problem for \tilde{A}_\pm^0 in (4.1) has a solution $I \pm \tilde{T}$ for some $\tilde{T} = \tilde{T}^*$ with $\nu_-(I - \tilde{T}^2) = \kappa$ if and only if the following condition is satisfied:*

$$\nu_-(I - \tilde{T}_{11}^2) = \nu_-(I - \tilde{T}_1^* \tilde{T}_1). \quad (4.2)$$

If this condition is satisfied then the following facts hold:

- (i) *The completion problems for \tilde{A}_\pm^0 in (4.1) have minimal solutions \tilde{A}_\pm .*
- (ii) *The operators $\tilde{T}_m := \tilde{A}_+ - I$ and $\tilde{T}_M := I - \tilde{A}_- \in \text{Ext}_{\tilde{T}_{11}, \kappa}(-1, 1)$.*
- (iii) *The operators \tilde{T}_m and \tilde{T}_M have the block form*

$$\begin{aligned} \tilde{T}_m &= \begin{pmatrix} \tilde{T}_{11} & D_{\tilde{T}_{11}} V^* \\ V D_{\tilde{T}_{11}} & -I + V(I - \tilde{T}_{11}) J_{11} V^* \end{pmatrix}, \\ \tilde{T}_M &= \begin{pmatrix} \tilde{T}_{11} & D_{\tilde{T}_{11}} V^* \\ V D_{\tilde{T}_{11}} & I - V(I + \tilde{T}_{11}) J_{11} V^* \end{pmatrix}, \end{aligned} \quad (4.3)$$

where $D_{\tilde{T}_{11}} := |I - \tilde{T}_{11}^2|^{1/2}$ and V is given by $V := \text{clos}(\tilde{T}_{21}D_{\tilde{T}_{11}}^{[-1]})$.

(iv) The operators \tilde{T}_m and \tilde{T}_M are extremal extensions of \tilde{T}_1 :

$$\tilde{T} \in \text{Ext}_{\tilde{T}_1, \kappa}(-1, 1) \text{ iff } \tilde{T} = \tilde{T}^* \in [\mathfrak{H}], \quad \tilde{T}_m \leq \tilde{T} \leq \tilde{T}_M.$$

(v) The operators \tilde{T}_m and \tilde{T}_M are connected via

$$(-\tilde{T})_m = -\tilde{T}_M, \quad (-\tilde{T})_M = -\tilde{T}_m.$$

For what follows it is convenient to reformulate the above theorem in a Kreĭn space setting. Consider the Kreĭn space (\mathfrak{H}, J) and a selfadjoint operator T in this space. Now the problem concerns selfadjoint operators $A_+ = I + T$ and $A_- = I - T$ in the Kreĭn space (\mathfrak{H}, J) that solve the completion problems

$$A_{\pm}^0 = \begin{pmatrix} I \pm T_{11} & \pm T_{21}^{[*]} \\ \pm T_{21} & * \end{pmatrix}, \quad (4.4)$$

under *minimal index conditions* $\nu_-(I + JT) = \nu_-(I + J_1 T_{11})$ and $\nu_-(I - JT) = \nu_-(I - J_1 T_{11})$, respectively. The set of solutions T to the problem (4.4) will be denoted by $\text{Ext}_{J_2 T_1, \kappa}(-1, 1)$.

Denote

$$T_1 = \begin{pmatrix} T_{11} \\ T_{21} \end{pmatrix} : (\mathfrak{H}_1, J_1) \rightarrow \begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix}, \quad (4.5)$$

so that T_1 is symmetric (nondensely defined) operator in the Kreĭn space $[(\mathfrak{H}_1, J_1)]$, i.e. $T_{11} = T_{11}^{[*]}$.

Theorem 4.2. *Let T_1 be a symmetric operator in a Kreĭn space sense as in (4.5) with $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ and $\nu_-(I - T_{11}^* T_{11}) = \kappa < \infty$, and let $J = \text{sign}(I - T_{11}^* T_{11})$. Then the completion problems for A_{\pm}^0 in (4.4) have a solution $I \pm T$ for some $T = T^{[*]}$ with $\nu_-(I - T^* T) = \kappa$ if and only if the following condition is satisfied:*

$$\nu_-(I - T_{11}^* T_{11}) = \nu_-(I - T_1^* T_1). \quad (4.6)$$

If this condition is satisfied then the following facts hold:

- (i) The completion problems for A_{\pm}^0 in (4.4) have "minimal" (J_2 -minimal) solutions A_{\pm} .
- (ii) The operators $T_m := A_+ - J$ and $T_M := J - A_- \in \text{Ext}_{J_2 T_1, \kappa}(-1, 1)$.
- (iii) The operators T_m and T_M have the block form

$$\begin{aligned} T_m &= \begin{pmatrix} T_{11} & J_1 D_{T_{11}} V^* \\ J_2 V D_{T_{11}} & -J_2 + J_2 V (I - J_1 T_{11}) J_{11} V^* \end{pmatrix}, \\ T_M &= \begin{pmatrix} T_{11} & J_1 D_{T_{11}} V^* \\ J_2 V D_{T_{11}} & J_2 - J_2 V (I + J_1 T_{11}) J_{11} V^* \end{pmatrix}, \end{aligned} \quad (4.7)$$

where $D_{T_{11}} := |I - T_{11}^* T_{11}|^{1/2}$ and V is given by $V := \text{clos}(J_2 T_{21} D_{T_{11}}^{[-1]})$.

(iv) The operators T_m and T_M are J_2 -extremal extensions of T_1 :

$$T \in \text{Ext}_{J_2 T_1, \kappa}(-1, 1) \text{ iff } T = T^{[*]} \in [(\mathfrak{H}, J)], \quad T_m \leq_{J_2} T \leq_{J_2} T_M.$$

(v) The operators T_m and T_M are connected via

$$(-T)_m = -T_M, \quad (-T)_M = -T_m.$$

Proof. The proof is obtained by systematic use of the equivalence that T is a selfadjoint operator in a Kreĭn space if and only if \tilde{T} is a selfadjoint in a Hilbert space. In particular, T gives solutions to the completion problems (4.4) if and only if \tilde{T} solves the completion problems (4.4). In view of

$$I - T_{11}^* T_{11} = I - T_{11}^* J J T_{11} = I - \tilde{T}_{11}^2,$$

we are getting formula (4.6) from (4.2). Then formula (4.7) follows by multiplying the operators in (4.3) by the fundamental symmetry. \square

5. Completion problem in a Pontryagin space

5.1. Defect operators and link operators

Let $(\mathfrak{H}, (\cdot, \cdot))$ be a Hilbert space and let J be a symmetry in \mathfrak{H} , i.e. $J = J^* = J^{-1}$, so that $(\mathfrak{H}, (J\cdot, \cdot))$, becomes a Pontryagin space. Then associate with $T \in [\mathfrak{H}]$ the corresponding defect and signature operators

$$D_T = |J - T^*JT|^{1/2}, \quad J_T = \text{sign}(J - T^*JT), \quad \mathfrak{D}_T = \overline{\text{ran}} D_T,$$

where the so-called defect subspace \mathfrak{D}_T can be considered as a Pontryagin space with the fundamental symmetry J_T . Similar notations are used with T^* :

$$D_{T^*} = |J - TJT^*|^{1/2}, \quad J_{T^*} = \text{sign}(J - TJT^*), \quad \mathfrak{D}_{T^*} = \overline{\text{ran}} D_{T^*}.$$

By definition $J_T D_T^2 = J - T^*JT$ and $J_T D_T = D_T J_T$ with analogous identities for D_{T^*} and J_{T^*} . In addition,

$$(J - T^*JT)JT^* = T^*J(J - TJT^*), \quad (J - TJT^*)JT = TJ(J - T^*JT).$$

Recall that $T \in [\mathfrak{H}]$ is said to be a J -contraction if $J - T^*JT \geq 0$, i.e. $\nu_-(J - T^*JT) = 0$. If, in addition, T^* is a J -contraction, T is termed as a J -bicontraction.

For the following consideration an indefinite version of the commutation relation of the form $TD_T = D_{T^*}T$ is needed; these involve so-called link operators introduced in [3, Section 4] (see also [5]).

Definition 5.1. There exist unique operators $L_T \in [\mathfrak{D}_T, \mathfrak{D}_{T^*}]$ and $L_{T^*} \in [\mathfrak{D}_{T^*}, \mathfrak{D}_T]$ such that

$$D_{T^*}L_T = TJ D_T \upharpoonright \mathfrak{D}_T, \quad D_T L_{T^*} = T^*J D_{T^*} \upharpoonright \mathfrak{D}_{T^*}; \quad (5.1)$$

in fact, $L_T = D_{T^*}^{[-1]}TJ D_T \upharpoonright \mathfrak{D}_T$ and $L_{T^*} = D_T^{[-1]}T^*J D_{T^*} \upharpoonright \mathfrak{D}_{T^*}$.

The following identities can be obtained with direct calculations; see [3, Section 4]:

$$\begin{aligned} L_T^* J_{T^*} \upharpoonright \mathfrak{D}_{T^*} &= J_T L_{T^*}; \\ (J_T - D_T J D_T) \upharpoonright \mathfrak{D}_T &= L_T^* J_{T^*} L_T; \\ (J_{T^*} - D_{T^*} J D_{T^*}) \upharpoonright \mathfrak{D}_{T^*} &= L_{T^*}^* J_T L_{T^*}. \end{aligned} \quad (5.2)$$

Now let T be selfadjoint in Pontryagin space (\mathfrak{H}, J) , i.e. $T^* = JTJ$. Then connections between D_{T^*} and D_T , J_{T^*} and J_T , L_{T^*} and L_T can be established.

Lemma 5.2. Assume that $T^* = JTJ$. Then $D_T = |I - T^2|^{1/2}$ and the following equalities hold:

$$D_{T^*} = J D_T J, \quad (5.3)$$

in particular,

$$\begin{aligned} \mathfrak{D}_{T^*} &= J \mathfrak{D}_T \text{ and } \mathfrak{D}_T = J \mathfrak{D}_{T^*}; \\ J_{T^*} &= J J_T J; \end{aligned} \quad (5.4)$$

$$L_{T^*} = J L_T J. \quad (5.5)$$

Proof. The defect operator of T can be calculated by the formula

$$D_T = ((I - (T^*)^2) J J (I - T^2))^{1/4} = ((I - (T^*)^2) (I - T^2))^{1/4}.$$

Then

$$\begin{aligned} D_{T^*} &= (J (I - (T^*)^2) (I - T^2) J)^{1/4} = J ((I - (T^*)^2) (I - T^2))^{1/4} J \\ &= J D_T J \end{aligned}$$

i.e. (5.3) holds. This implies

$$J \mathfrak{D}_{T^*} \subset \mathfrak{D}_T \text{ and } J \mathfrak{D}_T \subset \mathfrak{D}_{T^*}.$$

Hence from the last two formulas we get

$$\mathfrak{D}_{T^*} = J(J \mathfrak{D}_{T^*}) \subset J \mathfrak{D}_T \subset \mathfrak{D}_{T^*}$$

and similarly

$$\mathfrak{D}_T = J(J \mathfrak{D}_T) \subset J \mathfrak{D}_{T^*} \subset \mathfrak{D}_T.$$

The formula

$$\begin{aligned} J_T D_T^2 &= J - T^* J T = J(J - T J T^*) J = J J_{T^*} D_{T^*}^2 J = J J_{T^*} J D_T^2 J J \\ &= J J_{T^*} J D_T^2 \end{aligned}$$

yields the equation (5.4).

The relation (5.5) follows from

$$D_T L_{T^*} = T^* J D_{T^*} \restriction \mathfrak{D}_{T^*} = J T J D_T J \restriction \mathfrak{D}_{T^*} = J D_{T^*} L_T J = D_T J L_T J. \quad \square$$

5.2. Lemmas on negative indices of certain block operators

The first two lemmas are of preparatory nature for the last two lemmas, which are used for the proof of the main theorem.

Lemma 5.3. *Let $\begin{pmatrix} J & T \\ T & J \end{pmatrix} : \begin{pmatrix} \mathfrak{H} \\ \mathfrak{H} \end{pmatrix} \rightarrow \begin{pmatrix} \mathfrak{H} \\ \mathfrak{H} \end{pmatrix}$ be a selfadjoint operator in the Hilbert space $\mathfrak{H}^2 = \mathfrak{H} \oplus \mathfrak{H}$. Then*

$$\left| \begin{pmatrix} J & T \\ T & J \end{pmatrix} \right|^{1/2} = U \begin{pmatrix} |J+T|^{1/2} & 0 \\ 0 & |J-T|^{1/2} \end{pmatrix} U^*,$$

where $U = \frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix}$ is a unitary operator.

Proof. It is easy to check that

$$\begin{pmatrix} J & T \\ T & J \end{pmatrix} = U \begin{pmatrix} J+T & 0 \\ 0 & J-T \end{pmatrix} U^*. \quad (5.6)$$

Then by taking the modulus one gets

$$\left| \begin{pmatrix} J & T \\ T & J \end{pmatrix} \right|^2 = \left(\begin{pmatrix} J & T \\ T & J \end{pmatrix}^* \begin{pmatrix} J & T \\ T & J \end{pmatrix} \right) = U \begin{pmatrix} |J+T|^2 & 0 \\ 0 & |J-T|^2 \end{pmatrix} U^*.$$

The last step is to extract the square roots (twice) from the both sides of the equation:

$$\left| \begin{pmatrix} J & T \\ T & J \end{pmatrix} \right|^{1/2} = U \begin{pmatrix} |J+T|^{1/2} & 0 \\ 0 & |J-T|^{1/2} \end{pmatrix} U^*.$$

The right hand side can be written in this form because U is unitary. \square

Lemma 5.4. *Let $T = T^* \in \mathfrak{H}$ be a selfadjoint operator in a Hilbert space \mathfrak{H} and let $J = J^* = J^{-1}$ be a fundamental symmetry in \mathfrak{H} with $\nu_-(J) < \infty$. Then*

$$\nu_-(J - TJT) + \nu_-(J) = \nu_-(J - T) + \nu_-(J + T). \quad (5.7)$$

In particular, $\nu_-(J - TJT) < \infty$ if and only if $\nu_-(J \pm T) < \infty$.

Proof. Consider block operators $\begin{pmatrix} J & T \\ T & J \end{pmatrix}$ and $\begin{pmatrix} J+T & 0 \\ 0 & J-T \end{pmatrix}$. Equality (5.6) yields $\nu_- \begin{pmatrix} J & T \\ T & J \end{pmatrix} = \nu_- \begin{pmatrix} J+T & 0 \\ 0 & J-T \end{pmatrix}$. The negative index of $\begin{pmatrix} J+T & 0 \\ 0 & J-T \end{pmatrix}$ equals $\nu_-(J - T) + \nu_-(J + T)$ and the negative index of $\begin{pmatrix} J & T \\ T & J \end{pmatrix}$ is easy to find by using the equality

$$\begin{pmatrix} J & T \\ T & J \end{pmatrix} = \begin{pmatrix} I & 0 \\ TJ & I \end{pmatrix} \begin{pmatrix} J & 0 \\ 0 & J - TJT \end{pmatrix} \begin{pmatrix} I & JT \\ 0 & I \end{pmatrix}. \quad (5.8)$$

Then one gets (5.7). \square

Let $(\mathfrak{H}_i, (J_i \cdot, \cdot))$ ($i = 1, 2$) and $(\mathfrak{H}, (J \cdot, \cdot))$ be Pontryagin spaces, where $\mathfrak{H} = \mathfrak{H}_1 \oplus \mathfrak{H}_2$ and $J = \begin{pmatrix} J_1 & 0 \\ 0 & J_2 \end{pmatrix}$. Consider an operator $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ such that $\nu_-[I - T_{11}^2] = \kappa < \infty$; see (1.1). Denote $\tilde{T}_{11} = J_1 T_{11}$, then $\tilde{T}_{11} = \tilde{T}_{11}^*$ in the Hilbert space \mathfrak{H}_1 . Rewrite

$$\begin{aligned} \nu_-[I - T_{11}^2] &= \nu_-(J_1(I - T_{11}^2)) = \nu_-(J_1 - \tilde{T}_{11}J_1\tilde{T}_{11}) \\ &= \nu_-((J_1 - \tilde{T}_{11})J_1(J_1 + \tilde{T}_{11})). \end{aligned}$$

Furthermore, denote

$$\begin{aligned} J_+ &= \text{sign}(J_1(I - T_{11})) = \text{sign}(J_1 - \tilde{T}_{11}), \\ J_- &= \text{sign}(J_1(I + T_{11})) = \text{sign}(J_1 + \tilde{T}_{11}), \\ J_{11} &= \text{sign}(J_1(I - T_{11}^2)) \end{aligned} \quad (5.9)$$

and let $\kappa_+ = \nu_-[I - T_{11}]$ and $\kappa_- = \nu_-[I + T_{11}]$. Notice that $|I \mp T_{11}| = |J_1 \mp \tilde{T}_{11}|$ and one has polar decompositions

$$I \mp T_{11} = J_1 J_{\pm} |I \mp T_{11}|. \quad (5.10)$$

Lemma 5.5. *Let $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ and $T = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix} \in [(\mathfrak{H}, J)]$ be a selfadjoint extension of the operator T_{11} with $\nu_-[I \pm T_{11}] < \infty$ and $\nu_-(J) < \infty$. Then the following statements*

- (i) $\nu_-[I \pm T_{11}] = \nu_-[I \pm T]$;
- (ii) $\nu_-[I - T^2] = \nu_-[I - T_{11}^2] - \nu_-(J_2)$;
- (iii) $\text{ran } J_1 T_{21}^{[*]} \subset \text{ran } |I \pm T_{11}|^{1/2}$

are connected by the implications (i) \Leftrightarrow (ii) \Rightarrow (iii).

Proof. The Lemma can be formulated in an equivalent way for the Hilbert space operators: the block operator $\tilde{T} = JT = \begin{pmatrix} \tilde{T}_{11} & \tilde{T}_{12} \\ \tilde{T}_{21} & \tilde{T}_{22} \end{pmatrix}$ is a selfadjoint

extension of $\tilde{T}_{11} = \tilde{T}_{11}^* \in [\mathfrak{H}_1]$. Then the following statements

- (i') $\nu_-(J_1 \pm \tilde{T}_{11}) = \nu_-(J \pm \tilde{T})$
- (ii') $\nu_-(J - \tilde{T}J\tilde{T}) = \nu_-(J_1 - \tilde{T}_{11}J_1\tilde{T}_{11}) - \nu_-(J_2)$;
- (iii') $\text{ran } \tilde{T}_{12} \subset \text{ran } |J_1 \pm \tilde{T}_{11}|^{1/2}$

are connected by the implications (i') \Leftrightarrow (ii') \Rightarrow (iii').

Hence it's sufficient to prove this form of the Lemma.

Let us prove the equivalence (i') \Leftrightarrow (ii'). Condition (ii') is equivalent to

$$\nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} = \nu_- \begin{pmatrix} J & \tilde{T} \\ \tilde{T} & J \end{pmatrix}. \quad (5.11)$$

Indeed, in view of (5.8)

$$\nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} = \nu_-(J_1) + \nu_-(J_1 - \tilde{T}_{11}J_1\tilde{T}_{11})$$

and

$$\nu_- \begin{pmatrix} J & \tilde{T} \\ \tilde{T} & J \end{pmatrix} = \nu_-(J) + \nu_-(J - \tilde{T}J\tilde{T}) = \nu_-(J_1) + \nu_-(J_2) + \nu_-(J - \tilde{T}J\tilde{T}).$$

By using Lemma 5.4, equality (5.11) is equivalent to

$$\nu_-(J_1 - \tilde{T}_{11}) + \nu_-(J_1 + \tilde{T}_{11}) = \nu_-(J - \tilde{T}) + \nu_-(J + \tilde{T}). \quad (5.12)$$

Hence, (i') \Rightarrow (ii').

Because $\nu_-(J_1 \pm \tilde{T}_{11}) \leq \nu_-(J \pm \tilde{T})$, then (5.12) shows that (ii') \Rightarrow (i').

Now we prove implication (ii') \Rightarrow (iii'); the arguments here will be useful also for the proof of Lemma 5.6 below. Use a permutation to transform the matrix in the right hand side of (5.11):

$$\nu_- \begin{pmatrix} J & \tilde{T} \\ \tilde{T} & J \end{pmatrix} = \nu_- \begin{pmatrix} J_1 & 0 & \tilde{T}_{11} & \tilde{T}_{12} \\ 0 & J_2 & \tilde{T}_{21} & \tilde{T}_{22} \\ \tilde{T}_{11} & \tilde{T}_{12} & J_1 & 0 \\ \tilde{T}_{21} & \tilde{T}_{22} & 0 & J_2 \end{pmatrix} = \nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} & 0 & \tilde{T}_{12} \\ \tilde{T}_{11} & J_1 & \tilde{T}_{12} & 0 \\ 0 & \tilde{T}_{21} & J_2 & \tilde{T}_{22} \\ \tilde{T}_{21} & 0 & \tilde{T}_{22} & J_2 \end{pmatrix}.$$

Then condition (5.11) implies to the condition

$$\text{ran } \begin{pmatrix} 0 & \tilde{T}_{12} \\ \tilde{T}_{12} & 0 \end{pmatrix} \subset \text{ran } \left| \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} \right|^{1/2};$$

(see Theorem 2.1). By Lemma 5.3 the last inclusion can be rewritten as

$$\operatorname{ran} \begin{pmatrix} 0 & \tilde{T}_{12} \\ \tilde{T}_{12} & 0 \end{pmatrix} \subset \operatorname{ran} U \begin{pmatrix} |J_1 + \tilde{T}_{11}|^{1/2} & 0 \\ 0 & |J_1 - \tilde{T}_{11}|^{1/2} \end{pmatrix} U^*,$$

where $U = \frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix}$ is a unitary operator. This inclusion is equivalent to

$$\begin{aligned} \operatorname{ran} U^* \begin{pmatrix} 0 & \tilde{T}_{12} \\ \tilde{T}_{12} & 0 \end{pmatrix} U &= \operatorname{ran} \begin{pmatrix} \tilde{T}_{12} & 0 \\ 0 & -\tilde{T}_{12} \end{pmatrix} \\ &\subset \operatorname{ran} \begin{pmatrix} |J_1 + \tilde{T}_{11}|^{1/2} & 0 \\ 0 & |J_1 - \tilde{T}_{11}|^{1/2} \end{pmatrix} \end{aligned}$$

and clearly this is equivalent to condition (iii').

Note that if \tilde{T}_{11} has a selfadjoint extension \tilde{T} satisfying (i'). Then by applying Theorem 2.1 (or [5, Theorem 1]) it yields (iii'). \square

Lemma 5.6. *Let $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ be an operator and let*

$$T_1 = \begin{pmatrix} T_{11} \\ T_{21} \end{pmatrix} : (\mathfrak{H}_1, J_1) \rightarrow \begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix}$$

be an extension of T_{11} with $\nu_-[I - T_{11}^2] < \infty$, $\nu_-(J_1) < \infty$, and $\nu_-(J_2) < \infty$. Then for the conditions

- (i) $\nu_-[I_1 - T_{11}^2] = \nu_-[I_1 - T_1^{[*]}T_1] + \nu_-(J_2)$;
- (ii) $\operatorname{ran} J_1 T_{21}^{[*]} \subset \operatorname{ran} |I - T_{11}^2|^{1/2}$;
- (iii) $\operatorname{ran} J_1 T_{21}^{[*]} \subset \operatorname{ran} |I \pm T_{11}|^{1/2}$

the implications (i) \Rightarrow (ii) and (i) \Rightarrow (iii) hold.

Proof. First we prove that (i) \Rightarrow (ii). In fact, this follows from Theorem 3.1 by taking $A = I - T_{11}^2$ and $B = T_{21}$.

A proof of (i) \Rightarrow (iii) is quite similar to the proof used in Lemma 5.5. Statement (i) is equivalent the following equation:

$$\nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} = \nu_- \begin{pmatrix} J & \tilde{T}_1 \\ \tilde{T}_1^* & J_1 \end{pmatrix}.$$

Indeed,

$$\begin{aligned} \nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} &= \nu_- \begin{pmatrix} J_1 & 0 \\ 0 & J_1 - \tilde{T}_{11}J_1\tilde{T}_{11} \end{pmatrix} \\ &= \nu_-(J_1 - \tilde{T}_{11}J_1\tilde{T}_{11}) + \nu_-(J_1) < \infty \end{aligned}$$

and

$$\begin{aligned} \nu_- \begin{pmatrix} J & \tilde{T}_1 \\ \tilde{T}_1^* & J_1 \end{pmatrix} &= \nu_- \begin{pmatrix} J & 0 \\ 0 & J_1 - \tilde{T}_1^*J\tilde{T}_1 \end{pmatrix} \\ &= \nu_-(J_1 - \tilde{T}_{11}J_1\tilde{T}_{11} - \tilde{T}_{21}^*J_2\tilde{T}_{21}) + \nu_-(J_1) + \nu_-(J_2). \end{aligned}$$

Due to (i) the right hand sides coincide and then the left hand sides coincide as well.

Now let us permute the matrix in the latter equation.

$$\nu_- \begin{pmatrix} J & \tilde{T}_1 \\ \tilde{T}_1^* & J_1 \end{pmatrix} = \nu_- \begin{pmatrix} J_1 & 0 & \tilde{T}_{11} \\ 0 & J_2 & \tilde{T}_{21} \\ \tilde{T}_{11} & \tilde{T}_{21}^* & J_1 \end{pmatrix} = \nu_- \begin{pmatrix} J_1 & \tilde{T}_{11} & 0 \\ \tilde{T}_{11} & J_1 & \tilde{T}_{21}^* \\ 0 & \tilde{T}_{21} & J_2 \end{pmatrix}.$$

It follows from [5, Theorem 1] that the condition (i) implies the condition

$$\begin{aligned} \operatorname{ran} \begin{pmatrix} 0 \\ \tilde{T}_{21}^* \end{pmatrix} &\subset \operatorname{ran} \left| \begin{pmatrix} J_1 & \tilde{T}_{11} \\ \tilde{T}_{11} & J_1 \end{pmatrix} \right|^{1/2} \\ &= \operatorname{ran} U \begin{pmatrix} |J_1 + \tilde{T}_{11}|^{1/2} & 0 \\ 0 & |J_1 - \tilde{T}_{11}|^{1/2} \end{pmatrix} U^*, \end{aligned}$$

where $U = \frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix}$ is a unitary operator (see Lemma 5.3). Then, equivalently,

$$\operatorname{ran} \tilde{T}_{21}^* \subset \operatorname{ran} |J_1 \pm \tilde{T}_{11}|^{1/2}. \quad \square$$

5.3. Contractive extensions of contractions with minimal negative indices

Following to [5, 12, 14] we consider the problem of existence and a description of selfadjoint operators T in the Pontryagin space $\begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix}$ such that $A_+ = I + T$ and $A_- = I - T$ solve the corresponding completion problems

$$A_{\pm}^0 = \begin{pmatrix} I \pm T_{11} & \pm T_{21}^{[*]} \\ \pm T_{21} & * \end{pmatrix}, \quad (5.13)$$

under *minimal index conditions* $\nu_-[I + T] = \nu_-[I + T_{11}]$, $\nu_-[I - T] = \nu_-[I - T_{11}]$, respectively. Observe, that by Lemma 5.5 the two minimal index conditions above are equivalent to single condition $\nu_-[I - T^2] = \nu_-[I - T_{11}^2] - \nu_-(J_2)$.

It is clear from Theorem 2.1 that the conditions $\operatorname{ran} J_1 T_{21}^{[*]} \subset \operatorname{ran} |I - T_{11}|^{1/2}$ and $\operatorname{ran} J_1 T_{21}^{[*]} \subset \operatorname{ran} |I + T_{11}|^{1/2}$ are necessary for the existence of solutions; however as noted already in [5] they are not sufficient even in the Hilbert space setting.

The next theorem gives a general solvability criterion for the completion problem (5.13) and describes all solutions to this problem. As in the definite case, there are minimal solutions A_+ and A_- which are connected to two extreme selfadjoint extensions T of

$$T_1 = \begin{pmatrix} T_{11} \\ T_{21} \end{pmatrix} : (\mathfrak{H}_1, J_1) \rightarrow \begin{pmatrix} (\mathfrak{H}_1, J_1) \\ (\mathfrak{H}_2, J_2) \end{pmatrix}, \quad (5.14)$$

now with finite negative index $\nu_-[I - T^2] = \nu_-[I - T_{11}^2] - \nu_-(J_2) > 0$. The set of solutions T to the problem (5.13) will be denoted by $\operatorname{Ext}_{T_1, \kappa}(-1, 1)_{J_2}$.

Theorem 5.7. *Let T_1 be a symmetric operator as in (5.14) with $T_{11} = T_{11}^{[*]} \in [(\mathfrak{H}_1, J_1)]$ and $\nu_-[I - T_{11}^2] = \kappa < \infty$, and let $J_{T_{11}} = \text{sign}(J_1(I - T_{11}^2))$. Then the completion problem for A_{\pm}^0 in (5.13) has a solution $I \pm T$ for some $T = T^{[*]}$ with $\nu_-[I - T^2] = \kappa - \nu_-(J_2)$ if and only if the following condition is satisfied:*

$$\nu_-[I - T_{11}^2] = \nu_-[I - T_1^{[*]}T_1] + \nu_-(J_2). \quad (5.15)$$

If this condition is satisfied then the following facts hold:

- (i) *The completion problems for A_{\pm}^0 in (5.13) have "minimal" solutions A_{\pm} (for the partial ordering introduced in the first section).*
- (ii) *The operators $T_m := A_+ - I$ and $T_M := I - A_- \in \text{Ext}_{T_1, \kappa}(-1, 1)_{J_2}$.*
- (iii) *The operators T_m and T_M have the block form*

$$\begin{aligned} T_m &= \begin{pmatrix} T_{11} & J_1 D_{T_{11}} V^* \\ J_2 V D_{T_{11}} & -I + J_2 V(I - L_T^* J_1) J_{11} V^* \end{pmatrix}, \\ T_M &= \begin{pmatrix} T_{11} & J_1 D_{T_{11}} V^* \\ J_2 V D_{T_{11}} & I - J_2 V(I + L_T^* J_1) J_{11} V^* \end{pmatrix}, \end{aligned} \quad (5.16)$$

where $D_{T_{11}} := |I - T_{11}^2|^{1/2}$ and V is given by $V := \text{clos}(J_2 T_{21} D_{T_{11}}^{[-1]})$.

- (iv) *The operators T_m and T_M are "extremal" extensions of T_1 :*

$$T \in \text{Ext}_{T_1, \kappa}(-1, 1)_{J_2} \text{ iff } T = T^{[*]} \in [(\mathfrak{H}, J)], \quad T_m \leq_{J_2} T \leq_{J_2} T_M. \quad (5.17)$$

- (v) *The operators T_m and T_M are connected via*

$$(-T)_m = -T_M, \quad (-T)_M = -T_m. \quad (5.18)$$

Proof. It is easy to see by (3.1) that $\kappa = \nu_-[I - T_{11}^2] \leq \nu_-[I - T_1^{[*]}T_1] + \nu_-(J_2) \leq \nu_-[I - T^2] + \nu_-(J_2)$. Hence the condition $\nu_-[I - T^2] = \kappa - \nu_-(J_2)$ implies (5.15). The sufficiency of this condition is obtained when proving the assertions (i)–(iii) below.

(i) If the condition (5.15) is satisfied then by using Lemma 5.6 one gets the inclusions $\text{ran } J_1 T_{21}^{[*]} \subset \text{ran } |I \pm T_{11}|^{1/2}$, which by Theorem 2.1 means that each of the completion problems, A_{\pm}^0 in (5.13), is solvable. It follows that the operators

$$S_- = |I + T_{11}|^{[-1/2]} J_1 T_{21}^{[*]}, \quad S_+ = |I - T_{11}|^{[-1/2]} J_1 T_{21}^{[*]} \quad (5.19)$$

are well defined and they provide the minimal solutions A_{\pm} to the completion problems for A_{\pm}^0 in (5.13).

(ii) & (iii) By Lemma 5.6 the inclusion $\text{ran } J_1 T_{21}^{[*]} \subset \text{ran } |I - T_{11}^2|^{1/2}$ holds. This inclusion is equivalent to the existence of a (unique) bounded operator $V^* = D_{T_{11}}^{[-1]} J_1 T_{21}^{[*]}$ with $\ker V \supset \ker D_{T_{11}}$, such that $J_1 T_{21}^{[*]} = D_{T_{11}} V^*$. The operators $T_m := A_+ - I$ and $T_M := I - A_-$ (see proof of (i)) by using (5.1), (5.2), and 5.2 can be now rewritten as in (5.16). Indeed, observe that (see

Theorem 2.1, (5.9), and (5.10))

$$\begin{aligned}
 J_2 S_-^* J_- S_- &= J_2 V D_{T_{11}} |I + T_{11}|^{[-1/2]} J_- |I + T_{11}|^{[-1/2]} D_{T_{11}} V^* \\
 &= J_2 V D_{T_{11}} (J_1 (I + T_{11}))^{[-1]} D_{T_{11}} V^* \\
 &= J_2 V D_{T_{11}} D_{T_{11}}^{[-1]} (I + L_{T_{11}}^* J_1)^{[-1]} D_{T_{11}} J_1 D_{T_{11}} V^* \\
 &= J_2 V (I + L_{T_{11}}^* J_1)^{[-1]} (J_{11} - L_{T_{11}}^* J_{T_{11}}^* L_{T_{11}}) V^* \\
 &= J_2 V (I + L_{T_{11}}^* J_1)^{[-1]} (J_{11} - (L_{T_{11}}^* J_1)^2 J_{11}) V^* \\
 &= J_2 V (I + L_{T_{11}}^* J_1)^{[-1]} (I + L_{T_{11}}^* J_1) (I - L_{T_{11}}^* J_1) J_{11} V^* \\
 &= J_2 V (I - L_{T_{11}}^* J_1) J_{11} V^*,
 \end{aligned}$$

where the third equality follows from (5.1) and the fourth from (5.2).

And similarly for

$$\begin{aligned}
 J_2 S_+^* J_+ S_+ &= J_2 V D_{T_{11}} |I - T_{11}|^{[-1/2]} J_+ |I - T_{11}|^{[-1/2]} D_{T_{11}} V^* \\
 &= J_2 V D_{T_{11}} (J_1 (I - T_{11}))^{[-1]} D_{T_{11}} V^* \\
 &= J_2 V D_{T_{11}} D_{T_{11}}^{[-1]} (I - L_{T_{11}}^* J_1)^{[-1]} D_{T_{11}} J_1 D_{T_{11}} V^* \\
 &= J_2 V (I - L_{T_{11}}^* J_1)^{[-1]} (J_{11} - L_{T_{11}}^* J_{T_{11}}^* L_{T_{11}}) V^* \\
 &= J_2 V (I - L_{T_{11}}^* J_1)^{[-1]} (J_{11} - (L_{T_{11}}^* J_1)^2 J_{11}) V^* \\
 &= J_2 V (I - L_{T_{11}}^* J_1)^{[-1]} (I - L_{T_{11}}^* J_1) (I + L_{T_{11}}^* J_1) J_{11} V^* \\
 &= J_2 V (I + L_{T_{11}}^* J_1) J_{11} V^*,
 \end{aligned}$$

which implies the representations for T_m and T_M in (5.16). Clearly, T_m and T_M are selfadjoint extensions of T_1 , which satisfy the equalities

$$\nu_-[I + T_m] = \kappa_-, \quad \nu_-[I - T_M] = \kappa_+.$$

Moreover, it follows from (5.16) that

$$T_M - T_m = \begin{pmatrix} 0 & 0 \\ 0 & 2(I - J_2 V J_{11} V^*) \end{pmatrix}. \quad (5.20)$$

Now the assumption (5.15) will be used again. Since $\nu_-[I - T_1^{[*]} T_1] = \nu_-[I - T_{11}^2] - \nu_-(J_2)$ and $T_{21} = J_2 V D_{T_{11}}$ it follows from Theorem 3.1 that $V^* \in [\mathfrak{H}_2, \mathfrak{D}_{T_{11}}]$ is J -contractive: $J_2 - V J_{11} V^* \geq 0$. Therefore, (5.20) shows that $T_M \geq_{J_2} T_m$ and $I + T_M \geq_{J_2} I + T_m$ and hence, in addition to $I + T_m$, also $I + T_M$ is a solution to the problem A_+^0 and, in particular, $\nu_-[I + T_M] = \kappa_- = \nu_-[I + T_m]$. Similarly, $I - T_M \leq_{J_2} I - T_m$ which implies that $I - T_m$ is also a solution to the problem A_-^0 , in particular, $\nu_-[I - T_m] = \kappa_+ = \nu_-[I - T_M]$. Now by applying Lemma 5.5 we get

$$\begin{aligned}
 \nu_-[I - T_m^2] &= \kappa - \nu_-(J_2), \\
 \nu_-[I - T_M^2] &= \kappa - \nu_-(J_2).
 \end{aligned}$$

Therefore, $T_m, T_M \in \text{Ext}_{T_1, \kappa}(-1, 1)_{J_2}$ which in particular proves that the condition (5.15) is sufficient for solvability of the completion problem (5.13).

(iv) Observe, that $T \in \text{Ext}_{T_1, \kappa}(-1, 1)_{J_2}$ if and only if $T = T^{[*]} \supset T_1$ and $\nu_-[I \pm T] = \kappa_{\mp}$. By Theorem 2.1 this is equivalent to

$$J_2 S_-^* J_- S_- - I \leq_{J_2} T_{22} \leq_{J_2} I - J_2 S_+^* J_+ S_+. \quad (5.21)$$

The inequalities (5.21) are equivalent to (5.17).

(v) The relations (5.18) follow from (5.19) and (5.16). □

Acknowledgements. The author thanks his supervisor Seppo Hassi for several detailed discussions on the results of this paper.

References

- [1] Antezana, J., Corach, G., and Stojanoff, D., Bilateral shorted operators and parallel sums. *Linear Algebra Appl.* **414** (2006), 570–588.
- [2] Arsene, Gr. and Gheondea, A., Completing Matrix Contractions, *J. Operator Theory*, **7** (1982), 179–189.
- [3] Arsene, Gr., Constantinescu, T., Gheondea, A., Lifting of Operators and Prescribed Numbers of Negative Squares, *Michigan Math. J.*, **34** (1987), 201–216.
- [4] Azizov, T.Ya. and Iokhvidov, I.S., *Linear operators in spaces with indefinite metric*, John Wiley and Sons, New York, 1989.
- [5] Baidiuk, D., and Hassi, S., Completion, extension, factorization, and lifting of operators, Arxiv 2014, (to appear in *Math. Ann.*)
- [6] Bognár, J., *Indefinite Inner Product Space*, Springer-Verlag, Berlin, 1974.
- [7] Constantinescu, T. and Gheondea, A.: Minimal Signature of Lifting operators. I, *J. Operator Theory*, **22** (1989), 345–367.
- [8] Constantinescu, T. and Gheondea, A.: Minimal Signature of Lifting operators. II, *J. Funct. Anal.*, **103** (1992), 317–352.
- [9] Davis, Ch., Kahan, W.M., and Weinberger, H.F., Norm preserving dilations and their applications to optimal error bounds, *SIAM J. Numer. Anal.*, **19**, no. 3 (1982), 445–469.
- [10] Dritschel, M.A., A lifting theorem for bicontractions on Kreĭn spaces, *J. Funct. Anal.*, **89** (1990), 61–89.
- [11] Dritschel, M.A. and Rovnyak, J., Extension theorems for contraction operators on Kreĭn spaces. *Extension and interpolation of linear operators and matrix functions*, 221–305, *Oper. Theory Adv. Appl.*, **47**, Birkhäuser, Basel, 1990.
- [12] Hassi, S., Malamud, M.M., and de Snoo, H.S.V., On Kreĭn’s Extension Theory of Nonnegative Operators, *Math. Nachr.*, **274/275** (2004), 40–73.
- [13] Kato, T., *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, Heidelberg, 1995.
- [14] Kolmanovich, V.U. and Malamud, M.M., Extensions of Sectorial operators and dual pair of contractions, (Russian) Manuscript No 4428-85. Deposited at Vses. Nauchn-Issled, Inst. Nauchno-Tech. Informatsii, VINITI 19 04 85, Moscow, *R. ZH Mat* 10B1144, (1985), 1–57.
- [15] Kreĭn, M.G., On hermitian operators with defect indices $(1, 1)$, *Dokl. Akad. Nauk SSSR*, **43** (1944), 339–342.
- [16] Kreĭn, M.G., On resolvents of Hermitian operator with deficiency index (m, m) , *Dokl. Akad. Nauk SSSR*, **52** (1946), 657–660.
- [17] Kreĭn, M.G., Theory of Selfadjoint Extensions of Semibounded Operators and Its Applications, I, *Mat. Sb.* **20**, No.3 (1947), 431–498.
- [18] Langer, H. and Textorius, B., Extensions of a bounded Hermitian operator T preserving the numbers of negative eigenvalues of $I - T^*T$, Research Report LiTH-MAT-R-87-17, Department of Mathematics, Linköping University, (1977), 15 pp.
- [19] Shmul’yan, Yu. L., A Hellinger operator integral, (Russian) *Mat. Sb. (N.S.)*, **49**, No.91 (1959), 381–430.
- [20] Shmul’yan, Yu. L. and Yanovskaya, R.N., Blocks of a contractive operator matrix, *Izv. Vyssh. Uchebn. Zaved. Mat.*, No. 7 (1981), 72–75.

Dmytro Baidiuk
Department of Mathematics and Statistics
University of Vaasa
P.O. Box 700, 65101 Vaasa
Finland
e-mail: dbaidiuk@uwasa.fi