

Asymptotic Properties of Parameter Estimators in Vasicek Model Driven by Tempered Fractional Brownian Motion

Yuliya Mishura 

Taras Shevchenko National University of Kyiv
Mälardalen University

Kostiantyn Ralchenko 

Taras Shevchenko National University of Kyiv
University of Vaasa

Olena Dehtiar

Taras Shevchenko National University of Kyiv

Abstract

The paper focuses on the Vasicek model driven by a tempered fractional Brownian motion. We derive the asymptotic distributions of the least-squares estimators (based on continuous-time observations) for the unknown drift parameters. This work continues the investigation by Mishura and Ralchenko (*Fractal and Fractional*, 8(2:79), 2024), where these estimators were introduced and their strong consistency was proved.

Keywords: tempered fractional process, tempered fractional Vasicek model, parameter estimation, asymptotic distribution.

1. Introduction

The main goal of this paper is to establish the asymptotic distributions of the estimators of the drift parameters in the tempered fractional Vasicek model, or, in other words, in the Vasicek model involving tempered fractional Brownian motion (TFBM) of the first kind, as the driver. This tempered process was introduced and studied by Meerschaert and Sabzikar (2013, 2014). Concerning the estimators, we use the estimators constructed in Mishura and Ralchenko (2024), where we established their strong consistency. Moreover, we applied the main results about asymptotic normality of the drift parameter's estimators in the Vasicek model with the Gaussian driver of the unspecified form, but satisfying several assumptions, proved in Theorem 3.2 of Es-Sebaïy and Es-Sebaïy (2021). However, not all conditions of the specified theorem are satisfied for our model; therefore, direct application of results from

Es-Sebaiy and Es.Sebaiy (2021) was impossible, and we had to significantly modify the main proofs. More information about the relation between our assumptions and those of Es-Sebaiy and Es.Sebaiy (2021) is provided in Appendix A.

The tempered fractional Vasicek model is described by the following stochastic differential equation:

$$dY_t = (a + bY_t) dt + \sigma dB_{H,\lambda}(t), \quad t \geq 0, \quad Y_0 = y_0, \quad (1.1)$$

where $a \in \mathbb{R}$, $b > 0$, $\sigma > 0$, $y_0 \in \mathbb{R}$ are constants, and $B_{H,\lambda} = \{B_{H,\lambda}, t \geq 0\}$ is a tempered fractional Brownian motion introduced in Meerschaert and Sabzikar (2013).

We focus on the case where $b > 0$ and continue to investigate the asymptotic behavior of the least squares estimator of the unknown parameters (a, b) . In Mishura and Ralchenko (2024), the strong consistency of the estimator was proved. In the present paper, we determine its asymptotic distribution. The main result indicates that, similar to the non-ergodic fractional Vasicek model studied in Es-Sebaiy and Es.Sebaiy (2021), the estimator of a is asymptotically normal, whereas the estimator of b follows a Cauchy-type asymptotic distribution. However, in the model (1.1), the estimators of a and b are not asymptotically independent.

Our proofs rely on the asymptotic behavior of the variance of TFBM and the asymptotic growth with probability one of its sample paths. These properties were established in Azmoodeh, Mishura, and Sabzikar (2022) and Mishura and Ralchenko (2024), respectively.

Parameter estimation for a model similar to (1.1) but driven by fractional Brownian motion, known as the fractional Vasicek model, has been extensively studied for over 20 years. The case $a = 0$, known as the fractional Ornstein–Uhlenbeck process, has been particularly well-studied. Drift parameter estimation for this case began in 2002 with the maximum likelihood estimation (MLE) discussed in Kleptsyna and Le Breton (2002); Tudor and Viens (2007), and the asymptotic and exact distributions of the MLE were later investigated in Tanaka (2013, 2015).

Alternative approaches to drift parameter estimation for the fractional Ornstein–Uhlenbeck process are found in Belfadli, Es-Sebaiy, and Ouknine (2011); Hu and Nualart (2010); Hu, Nualart, and Zhou (2019); Kubilius, Mishura, Ralchenko, and Seleznev (2015). Since the asymptotic behavior of this process and the estimators is significantly affected by the sign of the drift parameter, hypothesis testing methods for it were developed in Kukush, Mishura, and Ralchenko (2017); Moers (2012). For a comprehensive survey on drift parameter estimation in fractional diffusion models, see Mishura and Ralchenko (2017), and for a detailed presentation, we refer to the book Kubilius, Mishura, and Ralchenko (2017).

Drift parameter estimation for Ornstein–Uhlenbeck processes driven by more general and related Gaussian processes was considered in Chen and Zhou (2021); El Machkouri, Es-Sebaiy, and Ouknine (2016); Shen, Yin, and Yan (2016); Mendy (2013); Mishura, Ralchenko, and Shklyar (2023). A similar problem for complex-valued fractional Ornstein–Uhlenbeck processes with fractional noise was investigated in Chen, Hu, and Wang (2017). In Shen and Yu (2019), the least squares estimator for the drift of Ornstein–Uhlenbeck processes with small fractional Lévy noise was constructed and studied.

In the general case of a fractional Vasicek model with two unknown drift parameters, the least squares and ergodic-type estimators were studied in Lohvinenko, Ralchenko, and Zhuchenko (2016); Xiao and Yu (2019a,b), while the corresponding MLEs were investigated in Lohvinenko and Ralchenko (2017, 2018, 2019); Tanaka, Xiao, and Yu (2020). In Li and Dong (2018), the least squares estimators of the Vasicek-type model driven by sub-fractional Brownian motion were studied. The same problem for the case of more general Gaussian noise (including fractional, sub-fractional, and bifractional Brownian motions) was investigated in Es-Sebaiy and Es.Sebaiy (2021). Least squares estimation of the drift parameters for the approximate fractional Vasicek process was investigated in Wang, Xiao, and Li (2023).

Several papers are devoted to the model (1.1) with non-Gaussian noises. In particular, drift parameter estimation for a Vasicek model driven by a Hermite process was studied in Nourdin

and Tran (2019); Vasicek-type models with Lévy processes were considered in Es-Sebaiy, Al-Foraih, and Alazemi (2021); Kawai (2013).

It is worth mentioning that the theory of parameter estimation for stochastic differential equations driven by a standard Wiener process, especially for classical Ornstein–Uhlenbeck and Vasicek models, is now well-developed. For comprehensive resources, see the books Bishwal (2008); Iacus (2008); Kutoyants (2004); Liptser and Shiryaev (2001). More recent results in this direction can be found in Kubilius *et al.* (2017) and the papers Jiang and Dong (2015); Jiang, Liu, and Zhou (2020); Prykhodko and Ralchenko (2024); Shimizu (2012); Tang and Chen (2009). Additionally, parameter estimation for the reflected Ornstein–Uhlenbeck process was studied in Zang and Zhang (2019), and for the threshold Ornstein–Uhlenbeck process in Hu and Xi (2022).

The structure of this paper is as follows. In the beginning of Section 2, we recall the definition and properties of the TFBM. Subsequently, we introduce the tempered fractional Vasicek model and the least-squares-type estimators for the drift parameters, and we formulate the main result concerning the asymptotic distributions of these estimators. All proofs are provided in Section 3. In Subsection 3.1, we express our estimators in terms of three Gaussian processes, which are three different integrals involving TFBM. Next, in Subsection 3.2, we determine the joint asymptotic distribution of these Gaussian processes. This enables us to derive the proof of the main theorem, which is detailed in Subsection 3.3. The paper is supplemented with two appendices. In Appendix A, we discuss the relation between our model and the conditions presented in Es-Sebaiy and Es-Sebaiy (2021). Appendix B provides brief information on the special functions that arise in the calculation of the asymptotic variances of the estimators.

2. Model description and main result

2.1. Tempered fractional Brownian motion

Let $W = \{W_x, x \in \mathbb{R}\}$ be a two-sided Wiener process, $H > 0$, $\lambda > 0$. According to Meerschaert and Sabzikar (2013), a tempered fractional Brownian motion (TFBM) is a zero mean stochastic process $B_{H,\lambda} = \{B_{H,\lambda}(t), t \geq 0\}$ defined by the following Wiener integral

$$B_{H,\lambda}(t) = \int_{\mathbb{R}} \left[\exp\{-\lambda(t-x)_+\} (t-x)_+^{H-\frac{1}{2}} - \exp\{-\lambda(-x)_+\} (-x)_+^{H-\frac{1}{2}} \right] dW_x.$$

Its covariance function has the following form Meerschaert and Sabzikar (2013)

$$\text{Cov}[B_{H,\lambda}(t), B_{H,\lambda}(s)] = \frac{1}{2} \left(C_t^2 t^{2H} + C_s^2 s^{2H} - C_{|t-s|}^2 |t-s|^{2H} \right), \quad (2.1)$$

with

$$C_t^2 = \frac{2\Gamma(2H)}{(2\lambda t)^{2H}} - \frac{2\Gamma(H + \frac{1}{2})}{\sqrt{\pi} (2\lambda t)^H} K_H(\lambda t), \quad (2.2)$$

where $K_\nu(z)$ is the modified Bessel function of the second kind, see Appendix B.

The variance function of TFBM with parameters $H > 0$ and $\lambda > 0$ satisfies

$$\lim_{t \rightarrow +\infty} \mathbf{E}[B_{H,\lambda}(t)]^2 = \lim_{t \rightarrow +\infty} C_t^2 t^{2H} = \frac{2\Gamma(2H)}{(2\lambda)^{2H}}, \quad (2.3)$$

see (Azmoodeh *et al.* 2022, Proposition 2.4).

Furthermore, it was proved in (Mishura and Ralchenko 2024, Theorem 3) that for any $\delta > 0$, there exists a non-negative random variable $\xi = \xi(\delta)$ such that for all $t > 0$

$$\sup_{s \in [0, t]} |B_{H,\lambda}(s)| \leq (t^\delta \vee 1) \xi \quad \text{a.s.}, \quad (2.4)$$

and there exist positive constants $C_1 = C_1(\delta)$ and $C_2 = C_2(\delta)$ such that for all $u > 0$

$$\mathbf{P}(\xi > u) \leq C_1 e^{-C_2 u^2}.$$

2.2. Parameter estimation in the tempered fractional Vasicek model

We focus in this paper on the drift parameter estimation for the tempered fractional Vasicek model, which is described by the following stochastic differential equation:

$$Y_t = y_0 + \int_0^t (a + bY_s) ds + \sigma B_{H,\lambda}(t), \quad t > 0, \quad Y_0 = y_0, \quad (2.5)$$

where $a \in \mathbb{R}$, $b > 0$, $\sigma > 0$, $y_0 \in \mathbb{R}$. The solution $Y = \{Y_t, t \geq 0\}$ is given explicitly by

$$Y_t = \left(y_0 + \frac{a}{b}\right) e^{bt} - \frac{a}{b} + \sigma \int_0^t e^{b(t-s)} dB_{H,\lambda}(s), \quad (2.6)$$

where the integral is defined by the integration by parts:

$$\int_0^t e^{b(t-s)} dB_{H,\lambda}(s) := B_{H,\lambda}(t) + b \int_0^t e^{b(t-s)} B_{H,\lambda}(s) ds. \quad (2.7)$$

Let us consider the estimation of unknown drift parameter $\theta = (a, b) \in \mathbb{R} \times (0, \infty)$ in the model (2.5). Following Mishura and Ralchenko (2024), we define the estimator $\hat{\theta}_T = (\hat{a}_T, \hat{b}_T)$ as follows

$$\hat{a}_T = \frac{(Y_T - y_0) \left(\int_0^T Y_t^2 dt - \frac{1}{2} (Y_T + y_0) \int_0^T Y_t dt \right)}{T \int_0^T Y_t^2 dt - \left(\int_0^T Y_t dt \right)^2}, \quad (2.8)$$

$$\hat{b}_T = \frac{(Y_T - y_0) \left(\frac{1}{2} T (Y_T + y_0) - \int_0^T Y_t dt \right)}{T \int_0^T Y_t^2 dt - \left(\int_0^T Y_t dt \right)^2}. \quad (2.9)$$

According to (Mishura and Ralchenko 2024, Theorem 6), (\hat{a}_T, \hat{b}_T) is a strongly consistent estimator of the parameter (a, b) as $T \rightarrow \infty$. The purpose of the present paper is to find asymptotic distributions of \hat{a}_T and \hat{b}_T . More precisely, we shall prove that \hat{a}_T is asymptotically normal, and \hat{b}_T has asymptotic Cauchy-type distribution.

2.3. Main result

Let us introduce the notations

$$\alpha_{H,\lambda}^2 = \frac{2\Gamma(2H)}{(2\lambda)^{2H}} \quad \text{and} \quad \beta_{H,\lambda,b}^2 = \frac{b}{2} \int_0^\infty \exp\{-bu\} C_u^2 u^{2H} du. \quad (2.10)$$

The following theorem is the main result of the paper.

Theorem 2.1. *The estimators \hat{a}_T and \hat{b}_T from (2.8) and (2.9), respectively, have the following asymptotic properties.*

(i) *The estimator \hat{a}_T is asymptotically normal:*

$$T(\hat{a}_T - a) \xrightarrow{d} \mathcal{N}\left(0, \sigma^2 \alpha_{H,\lambda}^2\right) \quad \text{as } T \rightarrow \infty. \quad (2.11)$$

(ii) *The estimator \hat{b}_T has asymptotic Cauchy-type distribution:*

$$e^{bT} (\hat{b}_T - b) \xrightarrow{d} \frac{\eta_1}{\eta_2}, \quad \text{as } T \rightarrow \infty,$$

where $\eta_1 \simeq \mathcal{N}(0, 4b^2 \sigma^2 \beta_{H,\lambda,b}^2)$ and $\eta_2 \simeq \mathcal{N}(y_0 + \frac{a}{b}, \sigma^2 \beta_{H,\lambda,b}^2)$ are independent normal random variables.

Remark 2.2 (Joint distribution of the estimators). Unlike the case of the Vasicek model driven by fractional Brownian motion (see (Es-Sebaïy and Es-Sebaïy 2021, Proposition 4.1)), the estimators \hat{a}_T and \hat{b}_T are *not* asymptotically independent. More precisely, the following convergence holds

$$\left(\begin{array}{c} T(\hat{a}_T - a) \\ e^{bT}(\hat{b}_T - b) \end{array} \right) \xrightarrow{d} \left(\begin{array}{c} b\sigma\xi_2 - \sigma\xi_3 \\ \frac{2b\sigma\xi_3}{y_0 + \frac{a}{b} + b\sigma\xi_1} \end{array} \right), \quad (2.12)$$

where the random vector (ξ_1, ξ_2, ξ_3) has a Gaussian distribution $\mathcal{N}(\mathbf{0}, \Sigma)$ with the covariance matrix Σ defined in Proposition 3.9 below. The normal random variables $\xi_1 \simeq \mathcal{N}(0, b^{-2}\beta_{H,\lambda,b}^2)$ and $\xi_3 \simeq \mathcal{N}(0, \beta_{H,\lambda,b}^2)$ are independent, and so are $\xi_2 \simeq \mathcal{N}(0, b^{-2}(\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2))$ and ξ_3 . However, there is a correlation between ξ_1 and ξ_2 , namely $\text{Cov}(\xi_1, \xi_2) = b^{-2}\beta_{H,\lambda,b}^2$.

Remark 2.3 (Representation for $\beta_{H,\lambda,b}^2$ via hypergeometric function). The constant $\beta_{H,\lambda,b}^2$ can be represented in an alternative form, which may be more suitable for its numerical computation. Using formula (2.2) for C_t , we can rewrite it in the following form

$$\begin{aligned} \beta_{H,\lambda,b}^2 &= \frac{b}{2} \int_0^\infty \exp\{-bt\} \left(\frac{2\Gamma(2H)}{(2\lambda)^{2H}} - \frac{2\Gamma(H + \frac{1}{2})t^H}{\sqrt{\pi}(2\lambda)^H} K_H(\lambda t) \right) dt \\ &= \frac{\Gamma(2H)}{(2\lambda)^{2H}} - \frac{b\Gamma(H + \frac{1}{2})}{\sqrt{\pi}(2\lambda)^H} \int_0^\infty \exp\{-bt\} t^H K_H(\lambda t) dt. \end{aligned} \quad (2.13)$$

Furthermore, by (Gradshteyn and Ryzhik 2007, formula 6.621-3),

$$\begin{aligned} \int_0^\infty \exp\{-bt\} t^H K_H(\lambda t) dt \\ = \frac{\sqrt{\pi}(2\lambda)^H}{(b + \lambda)^{2H+1}} \frac{\Gamma(2H + 1)}{\Gamma(H + \frac{3}{2})} {}_2F_1\left(2H + 1, H + \frac{1}{2}; H + \frac{3}{2}; \frac{b - \lambda}{b + \lambda}\right), \end{aligned} \quad (2.14)$$

where ${}_2F_1$ denotes the Gauss hypergeometric function, see Appendix B. Hence, combining (2.13)–(2.14) and using the relation $\Gamma(H + \frac{3}{2}) = (H + \frac{1}{2})\Gamma(H + \frac{1}{2})$, we arrive at

$$\begin{aligned} \beta_{H,\lambda,b}^2 &= \frac{\Gamma(2H)}{(2\lambda)^{2H}} - \frac{2b\Gamma(2H + 1)}{(b + \lambda)^{2H+1}(2H + 1)} {}_2F_1\left(2H + 1, H + \frac{1}{2}; H + \frac{3}{2}; \frac{b - \lambda}{b + \lambda}\right) \\ &= \frac{1}{2}\alpha_{H,\lambda}^2 - \frac{2b\Gamma(2H + 1)}{(b + \lambda)^{2H+1}(2H + 1)} {}_2F_1\left(2H + 1, H + \frac{1}{2}; H + \frac{3}{2}; \frac{b - \lambda}{b + \lambda}\right). \end{aligned} \quad (2.15)$$

3. Proofs

Let us introduce the following processes

$$Z_t := \int_0^t e^{-bs} B_{H,\lambda}(s) ds, \quad U_t = e^{-bt} \int_0^t e^{bs} B_{H,\lambda}(s) ds, \quad (3.1)$$

$$V_t := e^{-bt} \int_0^t e^{bs} dB_{H,\lambda}(s) = B_{H,\lambda}(t) - bU_t. \quad (3.2)$$

The proof of the main result will be conducted according to the following scheme. First, in subsection 3.1 we express $T(\hat{a}_T - a)$ and $e^{bT}(\hat{b}_T - b)$ via the processes Z, U and V and remainder terms, vanishing at infinity. Then in subsection 3.2 we find the joint asymptotic distribution of the Gaussian vector (Z_T, U_T, V_T) as $T \rightarrow \infty$. Finally, using these results along with the Slutsky theorem, we derive the limits in distribution for $T(\hat{a}_T - a)$ and $e^{bT}(\hat{b}_T - b)$ as $T \rightarrow \infty$ in subsection 3.3.

3.1. Representation of the estimators

Let us recall some well-known facts about the convergence of integrals involving tempered fractional Vasicek process Y . It was proved in [Mishura and Ralchenko \(2024\)](#) that the random variable

$$Z_\infty := \int_0^\infty e^{-bs} B_{H,\lambda}(s) ds \quad (3.3)$$

is well defined and the following convergences hold a.s. as $T \rightarrow \infty$:

$$e^{-bT} Y_T \rightarrow \zeta, \quad (3.4)$$

$$e^{-bT} \int_0^T Y_t dt \rightarrow \frac{1}{b} \zeta, \quad (3.5)$$

$$e^{-2bT} \int_0^T Y_t^2 dt \rightarrow \frac{1}{2b} \zeta^2, \quad (3.6)$$

$$T^{-1} e^{-bT} \int_0^T Y_t t dt \rightarrow \frac{1}{b} \zeta, \quad (3.7)$$

where

$$\zeta := y_0 + \frac{a}{b} + b\sigma Z_\infty, \quad (3.8)$$

see ([Mishura and Ralchenko 2024](#), Lemma 6).

Now we are ready to formulate and prove an auxiliary lemma, which is crucial for the proof of the main theorem. The lemma provides a representation of the estimator \hat{b}_T via the integrals Z_T and V_T defined in (3.1)–(3.2).

Lemma 3.1. *For all $T > 0$*

$$e^{bT} (\hat{b}_T - b) = \frac{\sigma V_T (y_0 + \frac{a}{b} + b\sigma Z_T)}{D_T} + R_T, \quad (3.9)$$

where

$$D_T := e^{-2bT} \left(\int_0^T Y_t^2 dt - \frac{1}{T} \left(\int_0^T Y_t dt \right)^2 \right) \rightarrow \frac{1}{2b} \zeta^2 \quad a.s., \text{ as } T \rightarrow \infty, \quad (3.10)$$

and

$$R_T \rightarrow 0 \quad a.s., \text{ as } T \rightarrow \infty. \quad (3.11)$$

Proof. By the definition (2.9) of the estimator \hat{b}_T ,

$$e^{bT} (\hat{b}_T - b) = \frac{F_T}{D_T}, \quad (3.12)$$

where the denominator D_T is defined by (3.10), and the the numerator F_T has the following form

$$\begin{aligned} F_T &= e^{-bT} (Y_T - y_0) \left(\frac{1}{2} (Y_T + y_0) - \frac{1}{T} \int_0^T Y_t dt \right) - b e^{-bT} \left(\int_0^T Y_t^2 dt - \frac{1}{T} \left(\int_0^T Y_t dt \right)^2 \right) \\ &= F_{1,T} + F_{2,T} + F_{3,T} + F_{4,T}, \end{aligned} \quad (3.13)$$

where

$$\begin{aligned} F_{1,T} &= \frac{1}{2} e^{-bT} (Y_T - y_0) (Y_T + y_0), & F_{2,T} &= -\frac{1}{T} e^{-bT} (Y_T - Y_0) \int_0^T Y_t dt, \\ F_{3,T} &= -b e^{-bT} \int_0^T Y_t^2 dt, & F_{4,T} &= \frac{b}{T} e^{-bT} \left(\int_0^T Y_t^2 dt \right)^2. \end{aligned}$$

Let us consider each of $F_{i,T}$ separately. Substituting the right-hand side of the equation (2.5) instead of the process Y , we rewrite the term $F_{1,T}$ as follows:

$$\begin{aligned} F_{1,T} &= \frac{1}{2}e^{-bT} \left(aT + b \int_0^T Y_t dt + \sigma B_{H,\lambda}(T) \right) \left(2y_0 + aT + b \int_0^T Y_t dt + \sigma B_{H,\lambda}(T) \right) \\ &= abTe^{-bT} \int_0^T Y_t dt + y_0 b e^{-bT} \int_0^T Y_t dt + \frac{1}{2}b^2 e^{-bT} \left(\int_0^T Y_t dt \right)^2 \\ &\quad + b\sigma e^{-bT} B_{H,\lambda}(T) \int_0^T Y_t dt + \frac{1}{2}e^{-bT} (aT + \sigma B_{H,\lambda}(T))(2y_0 + aT + \sigma B_{H,\lambda}(T)). \end{aligned} \quad (3.14)$$

Note that it follows from (2.6), (2.7), and (3.1) that the process Y has the following representation:

$$Y_T = \left(y_0 + \frac{a}{b} \right) e^{bT} - \frac{a}{b} + b\sigma e^{bT} Z_T + \sigma B_{H,\lambda}(T). \quad (3.15)$$

Moreover, expressing $b \int_0^T Y_t dt$ from the equation (2.5) and using (3.15), we get

$$b \int_0^T Y_t dt = Y_T - y_0 - aT - \sigma B_{H,\lambda}(T) = \left(y_0 + \frac{a}{b} \right) e^{bT} + b\sigma e^{bT} Z_T - y_0 - \frac{a}{b} - aT. \quad (3.16)$$

Now we insert (3.16) into the fourth term in the right-hand side of (3.14) and obtain

$$\begin{aligned} F_{1,T} &= abTe^{-bT} \int_0^T Y_t dt + y_0 b e^{-bT} \int_0^T Y_t dt + \sigma \left(y_0 + \frac{a}{b} \right) B_{H,\lambda}(T) \\ &\quad + b\sigma^2 Z_T B_{H,\lambda}(T) + R_{1,T}, \end{aligned} \quad (3.17)$$

where

$$R_{1,T} = \frac{1}{2}e^{-bT} (aT + \sigma B_{H,\lambda}(T))(2y_0 + aT + \sigma B_{H,\lambda}(T)) - \sigma e^{-bT} \left(y_0 + \frac{a}{b} + aT \right) B_{H,\lambda}(T).$$

In view of (2.4)

$$R_{1,T} \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty. \quad (3.18)$$

Let us consider $F_{2,T}$. By (2.5),

$$\begin{aligned} F_{2,T} &= -\frac{1}{T}e^{-bT} \left(aT + b \int_0^T Y_t dt + \sigma B_{H,\lambda}(T) \right) \int_0^T Y_t dt \\ &= -ae^{-bT} \int_0^T Y_t dt - \frac{b}{T}e^{-bT} \left(\int_0^T Y_t dt \right)^2 + R_{2,T}, \end{aligned} \quad (3.19)$$

where

$$R_{2,T} = -\frac{\sigma}{T}e^{-bT} B_{H,\lambda}(T) \int_0^T Y_t dt \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty, \quad (3.20)$$

due to (2.4) and (3.5).

Further, we transform $B_{3,T}$ using (2.5) as follows:

$$\begin{aligned} F_{3,T} &= -be^{-bT} \int_0^T Y_t \left(y_0 + at + b \int_0^t Y_s ds + \sigma B_{H,\lambda}(t) \right) dt \\ &= -by_0 e^{-bT} \int_0^T Y_t dt - abe^{-bT} \int_0^T tY_t dt - b^2 e^{-bT} \int_0^T Y_t \int_0^t Y_s ds dt \\ &\quad - b\sigma e^{-bT} \int_0^T Y_t B_{H,\lambda}(t) dt \\ &:= F_{31,T} + F_{32,T} + F_{33,T} + F_{34,T}. \end{aligned} \quad (3.21)$$

Integrating by parts and applying (3.16) we get

$$\begin{aligned}
F_{32,T} &= -abe^{-bT} \int_0^T t d \left(\int_0^t Y_s ds \right) = -abT e^{-bT} \int_0^T Y_t dt + abe^{-bT} \int_0^T \int_0^t Y_s ds dt \\
&= -abT e^{-bT} \int_0^T Y_t dt + ae^{-bt} \int_0^T (Y_t - y_0 - at - \sigma B_{H,\lambda}(t)) dt \\
&= -abT e^{-bT} \int_0^T Y_t dt + ae^{-bT} \int_0^T Y_t dt + R'_{3,T},
\end{aligned} \tag{3.22}$$

where

$$R'_{3,T} = -ae^{-bT} \int_0^T (y_0 + at + \sigma B_{H,\lambda}(t)) dt \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty \tag{3.23}$$

by (2.4).

Due to symmetry of the integrand, it is not hard to see that $\int_0^T \int_0^t Y_t Y_s ds dt = \frac{1}{2} (\int_0^T Y_t dt)^2$, whence

$$F_{33,T} = -\frac{b^2}{2} e^{-bT} \left(\int_0^T Y_t dt \right)^2. \tag{3.24}$$

In order to transform $F_{34,T}$, we use (3.16) and get

$$\begin{aligned}
F_{34,T} &= -b\sigma e^{-bT} \int_0^T B_{H,\lambda}(t) \left(\left(y_0 + \frac{a}{b} \right) e^{bt} - \frac{a}{b} + b\sigma e^{bt} Z_t + \sigma B_{H,\lambda}(t) \right) \\
&= -b\sigma \left(y_0 + \frac{a}{b} \right) e^{-bT} \int_0^T e^{bt} B_{H,\lambda}(t) dt + a\sigma e^{-bT} \int_0^T B_{H,\lambda}(t) dt \\
&\quad - b^2 \sigma^2 e^{-bT} \int_0^T e^{bt} B_{H,\lambda}(t) Z_t dt - b\sigma^2 e^{-bT} \int_0^T B_{H,\lambda}^2(t) dt.
\end{aligned}$$

Using integration by parts, we obtain

$$\begin{aligned}
\int_0^T e^{bt} B_{H,\lambda}(t) Z_t dt &= \int_0^T Z_t d \left(\int_0^t e^{bs} B_{H,\lambda}(s) ds \right) \\
&= Z_T \int_0^T e^{bt} B_{H,\lambda}(t) dt - \int_0^T \int_0^t e^{bs} B_{H,\lambda}(s) ds dZ_t.
\end{aligned}$$

Hence,

$$F_{34,T} = -b\sigma \left(y_0 + \frac{a}{b} \right) e^{bt} \int_0^T e^{bt} B_{H,\lambda}(t) dt - b^2 \sigma^2 e^{-bT} Z_T \int_0^T e^{bt} B_{H,\lambda}(t) dt + R''_{3,T},$$

where

$$\begin{aligned}
R''_{3,T} &= a\sigma e^{-bT} \int_0^T B_{H,\lambda}(t) dt - b^2 \sigma^2 e^{-bT} \int_0^T B_{H,\lambda}^2(t) dt \\
&\quad + b^2 \sigma^2 e^{-bT} \int_0^T \int_0^t e^{bs} B_{H,\lambda}(s) ds dZ_t.
\end{aligned} \tag{3.25}$$

Recall that by (3.1)–(3.2)

$$e^{-bT} \int_0^T e^{bt} B_{H,\lambda}(t) dt = U_T = \frac{1}{b} B_{H,\lambda}(T) - \frac{1}{b} V_T.$$

Therefore, we can rewrite $F_{34,T}$ as follows:

$$F_{34,T} = -\sigma \left(y_0 + \frac{a}{b} \right) B_{H,\lambda}(T) - b\sigma^2 B_{H,\lambda}(T) Z_T + \sigma V_T \left(y_0 + \frac{a}{b} + b\sigma Z_T \right) + R''_{3,T}. \tag{3.26}$$

Note that the first two terms in the right-hand side of (3.25) converge to zero a.s., as $T \rightarrow \infty$ due to (2.4). The last term in (3.25) also vanishes, because

$$\begin{aligned} \lim_{T \rightarrow \infty} e^{-bT} \int_0^T \int_0^t e^{bs} B_{H,\lambda}(s) ds dZ_t &= \lim_{T \rightarrow \infty} \frac{\int_0^T \int_0^t e^{bs} B_{H,\lambda}(s) ds e^{-bt} B_{H,\lambda}(t) dt}{e^{bT}} \\ &= \lim_{T \rightarrow \infty} \frac{\int_0^T e^{bs} B_{H,\lambda}(s) ds e^{-bT} B_{H,\lambda}(t)}{be^{bT}} = 0 \quad \text{a.s.} \end{aligned}$$

in view of (2.4).

Hence,

$$R''_{3,T} \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty. \tag{3.27}$$

Combining (3.13), (3.17), (3.19), (3.21), (3.22), (3.24), and (3.26), we arrive at

$$F_T = \sigma V_T (y_0 + \frac{a}{b} + b\sigma Z_T) + \tilde{R}_T, \tag{3.28}$$

where

$$\tilde{R}_T = R_{1,T} + R_{2,T} + R'_{3,T} + R''_{3,T} \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty, \tag{3.29}$$

by (3.18), (3.20), (3.23), and (3.27).

We complete the proof by inserting (3.28) into (3.12) and noticing that $R_T := \frac{\tilde{R}_T}{D_T} \rightarrow 0$ a.s., as $T \rightarrow \infty$ in view of (3.29) and (3.10). \square

In the next lemma, we express the estimator \hat{a}_T via the integrals U_T and V_T defined in (3.1)–(3.2). This representation also contains random variables P_T and Q_T , converging a.s. to the constants 1 and 0 respectively.

Lemma 3.2. *For all $T > 0$*

$$T(\hat{a}_T - a) = b\sigma U_T - \sigma V_T P_T + Q_T,$$

where

$$P_T := \frac{(y_0 + \frac{a}{b} + b\sigma Z_T) e^{-bT} \int_0^T Y_t dt}{D_T} - 1 \rightarrow 1 \quad \text{a.s., as } T \rightarrow \infty. \tag{3.30}$$

and

$$Q_T := -R_T e^{-bT} \int_0^T Y_t dt \rightarrow 0 \quad \text{a.s., as } T \rightarrow \infty. \tag{3.31}$$

Proof. Using (2.8) and (2.9) we rewrite $T\hat{a}_T$ as follows

$$\begin{aligned} T\hat{a}_T &= \frac{(Y_T - y_0) \left(T \int_0^T Y_t^2 dt - \left(\int_0^T Y_t dt \right)^2 + \left(\int_0^T Y_t dt \right)^2 - \frac{1}{2} T (Y_T + y_0) \int_0^T Y_t dt \right)}{T \int_0^T Y_t^2 dt - \left(\int_0^T Y_t dt \right)^2} \\ &= Y_T - y_0 - \hat{b}_T \int_0^T Y_t dt. \end{aligned}$$

Now expressing Y_T through (2.5), we get

$$T\hat{a}_T = Ta + \sigma B_{H,\lambda}(T) - (\hat{b}_T - b) \int_0^T Y_t dt. \tag{3.32}$$

Note that by (3.2), $B_{H,\lambda}(T) = bU_T + V_T$. Using this relation and the representation (3.9) we derive from (3.32) that

$$T(\hat{a}_T - a) = \sigma V_T + b\sigma U_T - \left(\frac{\sigma V_T (y_0 + \frac{a}{b} + b\sigma Z_T)}{D_T} + R_T \right) e^{-bT} \int_0^T Y_t dt$$

$$= b\sigma U_T - \sigma V_T P_T + Q_T.$$

Note that $y_0 + \frac{a}{b} + b\sigma Z_T \rightarrow \zeta$, $D_T \rightarrow \frac{1}{2b}\zeta^2$, $e^{-bT} \int_0^T Y_t dt \rightarrow \frac{1}{b}\zeta$ and $R_T \rightarrow 0$ a.s., as $T \rightarrow \infty$, by (3.8), (3.10), (3.5), and (3.11) respectively. This implies the convergences (3.30) and (3.31). \square

3.2. Asymptotic normality of (Z_T, U_T, V_T)

The purpose of this subsection is to find a joint asymptotic distribution of the integrals Z_T , U_T , and V_T as $T \rightarrow \infty$. This distribution is obviously Gaussian, since Z_T , U_T , and V_T are Gaussian processes. Therefore, it suffices to calculate the elements of the asymptotic covariance matrix. This will be done in the following series of lemmas. The limits contain the constants $\alpha_{H,\lambda}$ and $\beta_{H,\lambda,b}$ defined in Theorem 2.1.

Lemma 3.3. *The following convergence holds:*

$$\lim_{T \rightarrow \infty} \mathbf{E}Z_T^2 = \mathbf{E}Z_\infty^2 = \frac{\beta_{H,\lambda,b}^2}{b^2}. \quad (3.33)$$

Proof. Using the definition (3.3) of Z_∞ , and the formula (2.1) for the covariance function of TFBM, we may write

$$\begin{aligned} \mathbf{E}Z_\infty^2 &= \mathbf{E} \left(\int_0^\infty e^{-bt} B_{H,\lambda}(t) dt \right)^2 = \int_0^\infty \int_0^\infty e^{-bt-bs} \mathbf{E}[B_{H,\lambda}(t)B_{H,\lambda}(s)] dt ds \\ &= \frac{1}{2} \int_0^\infty \int_0^\infty e^{-bt-bs} \left(C_t^2 t^{2H} + C_s^2 s^{2H} - C_{|t-s|}^2 |t-s|^{2H} \right) ds dt \\ &= \int_0^\infty e^{-bs} ds \int_0^\infty e^{-bt} C_t^2 t^{2H} dt - \frac{1}{2} \int_0^\infty \int_0^\infty e^{-bt-bs} C_{|t-s|}^2 |t-s|^{2H} ds dt \\ &=: A_1 + A_2. \end{aligned} \quad (3.34)$$

Since $\int_0^\infty e^{-bs} ds = \frac{1}{b}$, we see that

$$A_1 = \frac{1}{b} \int_0^\infty e^{-bt} C_t^2 t^{2H} dt = \frac{2}{b^2} \beta_{H,\lambda,b}^2 \quad (3.35)$$

by definition of $\beta_{H,\lambda,b}$, see (2.10).

Let us consider A_2 . Due to the symmetry of the integrand, we have

$$A_2 = - \int_0^\infty \int_0^t e^{-bt-bs} C_{t-s}^2 (t-s)^{2H} ds dt = - \int_0^\infty \int_0^t e^{-2bt+bu} C_u^2 u^{2H} du dt,$$

where we have used the substitution $u = t - s$ in the inner integral. Changing the order of integration and integrating w.r.t. t , we then get

$$A_2 = - \int_0^\infty \left(\int_u^\infty e^{-2bt} dt \right) e^{bu} C_u^2 u^{2H} du = - \frac{1}{2b} \int_0^\infty e^{-bu} C_u^2 u^{2H} du = - \frac{1}{b^2} \beta_{H,\lambda,b}^2. \quad (3.36)$$

Combining (3.34)–(3.36), we obtain (3.33). \square

Lemma 3.4. *The following convergence holds:*

$$\lim_{T \rightarrow \infty} \mathbf{E}U_T^2 = \frac{1}{b^2} \left(\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2 \right). \quad (3.37)$$

Proof. Using the definition (3.1) of U_T and the formula (2.1) for the covariance function of TFBM, we have

$$\mathbf{E}U_T^2 = \mathbf{E} \left(\exp\{-bT\} \int_0^T \exp\{bs\} B_{H,\lambda}(s) ds \right)^2$$

$$\begin{aligned}
 &= \frac{1}{2} \exp\{-2bT\} \int_0^T \int_0^T \exp\{b(s+t)\} \left[C_t^2 t^{2H} + C_s^2 s^{2H} - C_{|t-s|}^2 |t-s|^{2H} \right] ds dt \\
 &= \exp\{-2bT\} \int_0^T \exp\{bs\} ds \int_0^T \exp\{bt\} C_t^2 t^{2H} dt \\
 &\quad - \frac{1}{2} \exp\{-2bT\} \int_0^T \int_0^T \exp\{b(s+t)\} C_{|t-s|}^2 |t-s|^{2H} ds dt \\
 &=: B_{1,T} + B_{2,T}.
 \end{aligned} \tag{3.38}$$

By the l'Hôpital rule and (2.3), we have

$$\lim_{T \rightarrow \infty} \frac{\int_0^T e^{bt} C_t^2 t^{2H} dt}{e^{bT}} = \lim_{T \rightarrow \infty} \frac{e^{bT} C_T^2 T^{2H}}{be^{bT}} = \frac{\alpha_{H,\lambda}^2}{b}, \tag{3.39}$$

whence

$$\lim_{T \rightarrow \infty} B_{1,T} = \lim_{T \rightarrow \infty} \frac{1 - e^{-bT}}{b} \cdot \frac{\int_0^T e^{bt} C_t^2 t^{2H} dt}{e^{bT}} = \frac{\alpha_{H,\lambda}^2}{b^2}. \tag{3.40}$$

Taking into account the symmetry of the integrand, we can represent $B_{2,T}$ in the following form:

$$\begin{aligned}
 B_{2,T} &= -\exp\{-2bT\} \int_0^T \int_0^t \exp\{b(s+t)\} C_{t-s}^2 (t-s)^{2H} ds dt \\
 &= -\exp\{-2bT\} \int_0^T \int_0^t \exp\{b(2t-u)\} C_u^2 u^{2H} du dt,
 \end{aligned}$$

where we have used the substitution $u = t - s$ in the inner integral. Changing the order of integration and integrating w.r.t. t , we obtain

$$\begin{aligned}
 B_{2,T} &= -\exp\{-2bT\} \int_0^T \exp\{-bu\} C_u^2 u^{2H} \int_u^T \exp\{2bt\} dt du \\
 &= -\frac{1}{2b} \int_0^T \exp\{-bu\} C_u^2 u^{2H} du + \frac{1}{2b} \exp\{-2bT\} \int_0^T \exp\{bu\} C_u^2 u^{2H} du.
 \end{aligned}$$

Note that the last term in the right-hand side of the above equality tends to zero due to (3.39). Therefore

$$\lim_{T \rightarrow \infty} B_{2,T} = -\frac{1}{2b} \int_0^\infty \exp\{-bu\} C_u^2 u^{2H} du = -\frac{1}{b^2} \beta_{H,\lambda,b}^2, \tag{3.41}$$

by the definition of $\beta_{H,\lambda,b}$, see (2.10). Combining (3.38), (3.40), and (3.41), we conclude the proof. \square

Lemma 3.5. *The following convergence holds:*

$$\lim_{T \rightarrow \infty} \mathbf{E}V_T^2 = \beta_{H,\lambda,b}^2. \tag{3.42}$$

Proof. Using (3.2), we represent the left-hand side of (3.42) in the following form

$$\mathbf{E}V_T^2 = \mathbf{E} (B_{H,\lambda}(T) - bU_T)^2 = \mathbf{E}B_{H,\lambda}^2(T) + b^2\mathbf{E}U_T^2 - 2b\mathbf{E} [B_{H,\lambda}(T)U_T]. \tag{3.43}$$

Next, we transform the third term in the right-hand side of (3.43) as follows

$$\begin{aligned}
 \mathbf{E} [B_{H,\lambda}(T)U_T] &= \exp\{-bT\} \int_0^T \exp\{bs\} \mathbf{E}[B_{H,\lambda}(T)B_{H,\lambda}(s)] ds \\
 &= \frac{1}{2} \exp\{-bT\} \int_0^T \exp\{bs\} \left[C_T^2 T^{2H} + C_s^2 s^{2H} - C_{T-s}^2 (T-s)^{2H} \right] ds \\
 &= \frac{1}{2} \exp\{-bT\} C_T^2 T^{2H} \int_0^T \exp\{bs\} ds + \frac{1}{2} \exp\{-bT\} \int_0^T \exp\{bs\} C_s^2 s^{2H} ds
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2} \exp\{-bT\} \int_0^T \exp\{bs\} C_{T-s}^2 (T-s)^{2H} ds \\
& = \frac{1}{2b} C_T^2 T^{2H} - \frac{1}{2b} \exp\{-bT\} C_T^2 T^{2H} + \frac{1}{2} \exp\{-bT\} \int_0^T \exp\{bs\} C_s^2 s^{2H} ds \\
& \quad - \frac{1}{2} \int_0^T \exp\{-bz\} C_z^2 z^{2H} dz. \tag{3.44}
\end{aligned}$$

Combining this expression with (3.43) and taking into account that $C_T^2 T^{2H} = \mathbf{E}[B_{H,\lambda}(T)^2]$ we arrive at

$$\begin{aligned}
\mathbf{E}V_T^2 & = b^2 \mathbf{E}U_T^2 + \exp\{-bT\} C_T^2 T^{2H} - b \exp\{-bT\} \int_0^T \exp\{bs\} C_s^2 s^{2H} ds \\
& \quad + b \int_0^T \exp\{-bz\} C_z^2 z^{2H} dz. \tag{3.45}
\end{aligned}$$

Note that according to (2.3) the second term in the right-hand side of (3.45) vanishes as $T \rightarrow \infty$, while the limits of other terms are already known, see (3.37), (3.39), and (2.10). Therefore, we arrive at

$$\lim_{T \rightarrow \infty} \mathbf{E}V_T^2 = \left(\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2 \right) - \alpha_{H,\lambda}^2 + 2\beta_{H,\lambda,b}^2 = \beta_{H,\lambda,b}^2. \quad \square$$

Lemma 3.6. *The following asymptotics holds*

$$\lim_{T \rightarrow \infty} \mathbf{E}[Z_T U_T] = \frac{\beta_{H,\lambda,b}^2}{b^2}.$$

Proof. Using formula (2.1) for the covariance function of TFBM, we get

$$\begin{aligned}
\mathbf{E}[Z_T U_T] & = e^{-bT} \mathbf{E} \left[\int_0^T e^{bt} B_{H,\lambda}(t) dt \int_0^T e^{-bs} B_{H,\lambda}(s) ds \right] \\
& = \frac{1}{2} e^{-bT} \int_0^T \int_0^T e^{bt-bs} \left(C_t^2 t^{2H} + C_s^2 s^{2H} - C_{|t-s|}^2 |t-s|^{2H} \right) ds dt \\
& = I_{1,T} + I_{2,T} + I_{3,T} + I_{4,T},
\end{aligned}$$

where

$$\begin{aligned}
I_{1,T} & = \frac{1}{2} e^{-bT} \int_0^T e^{bt} C_t^2 t^{2H} dt \int_0^T e^{-bs} ds, \\
I_{2,T} & = \frac{1}{2} e^{-bT} \int_0^T e^{-bs} C_s^2 s^{2H} ds \int_0^T e^{bt} dt, \\
I_{3,T} & = -\frac{1}{2} e^{-bT} \int_0^T \int_0^t e^{bt-bs} C_{t-s}^2 (t-s)^{2H} ds dt, \\
I_{4,T} & = -\frac{1}{2} e^{-bT} \int_0^T \int_t^T e^{bt-bs} C_{s-t}^2 (s-t)^{2H} ds dt.
\end{aligned}$$

By (3.39),

$$I_{1,T} = \frac{1}{2} e^{-bT} \cdot \frac{1 - e^{-bT}}{b} \int_0^T e^{bt} C_t^2 t^{2H} dt \rightarrow \frac{\alpha_{H,\lambda}^2}{2b^2}, \quad \text{as } T \rightarrow \infty.$$

Taking into account (2.10), we get

$$I_{2,T} = \frac{1}{2} e^{-bT} \cdot \frac{e^{-bT} - 1}{b} \int_0^T e^{-bs} C_s^2 s^{2H} ds \rightarrow \frac{\beta_{H,\lambda,b}^2}{b^2}, \quad \text{as } T \rightarrow \infty.$$

By l'Hopital's rule and (3.39), we have

$$I_{3,T} = - \lim_{T \rightarrow \infty} \frac{\int_0^T \int_0^t e^{bu} C_u^2 u^{2u} du}{2e^{bT}} = - \lim_{T \rightarrow \infty} \frac{\int_0^T e^{bu} C_u^2 u^{2u} du}{2be^{bT}} = - \frac{\alpha_{H,\lambda}^2}{2b^2}.$$

Changing the order of integration, we obtain

$$I_{4,T} = - \frac{1}{2} e^{-bT} \int_0^T \int_0^s e^{bt-bs} C_{s-t}^2 (s-t)^{2H} dt ds = - \frac{1}{2} e^{-bT} \int_0^T \int_0^s e^{-bu} C_u^2 u^{2H} du ds.$$

By l'Hopital's rule and (2.10),

$$\lim_{T \rightarrow \infty} I_{4,T} = - \lim_{T \rightarrow \infty} \frac{\int_0^T e^{-bu} C_u^2 u^{2u} du}{2be^{bT}} = 0.$$

Collecting all the limits completes the proof. □

Lemma 3.7. *The next value is asymptotically negligible:*

$$\lim_{T \rightarrow \infty} \mathbf{E} [Z_T V_T] = 0.$$

Proof. It follows from (3.2) that

$$\mathbf{E} [Z_T V_T] = \mathbf{E} [Z_T B_{H,\lambda}(T)] - b \mathbf{E} [Z_T U_T]. \tag{3.46}$$

By (2.3), (2.10), and (3.39), we obtain the next equalities

$$\begin{aligned} \mathbf{E} [Z_T B_{H,\lambda}(T)] &= \mathbf{E} \left[B_{H,\lambda}(T) \int_0^T e^{-bt} B_{H,\lambda}(t) dt \right] \\ &= \frac{1}{2} \int_0^T e^{-bT} \left(C_T^2 T^{2H} + C_t^2 t^{2H} - C_{T-t}^2 (T-t)^{2H} \right) dt \\ &= \frac{1}{2} C_T^2 T^{2H} \cdot \frac{1 - e^{-bT}}{b} + \frac{1}{2} \int_0^T e^{-bt} C_t^2 t^{2H} dt - \frac{1}{2} \int_0^T e^{-b(T-s)} C_s^2 s^{2H} ds \\ &\rightarrow \frac{\alpha_{H,\lambda}^2}{2b} + \frac{\beta_{H,\lambda,\beta}^2}{b} - \frac{\alpha_{H,\lambda}^2}{2b} = \frac{\beta_{H,\lambda,\beta}^2}{b}, \quad \text{as } T \rightarrow \infty. \end{aligned} \tag{3.47}$$

Combining (3.46), (3.47) and Lemma 3.6, we conclude the proof. □

Lemma 3.8. *The following value is also negligible:*

$$\lim_{T \rightarrow \infty} \mathbf{E} [U_T V_T] = 0.$$

Proof. From the representation (3.44) we derive using (2.3), (3.39), and (2.10) that

$$\lim_{T \rightarrow \infty} \mathbf{E} [B_{H,\lambda} U_T] = \frac{\alpha_{H,\lambda}^2}{2b} + 0 + \frac{\alpha_{H,\lambda}^2}{2b} - \frac{\beta_{H,\lambda,b}^2}{b} = \frac{\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2}{b} = b \lim_{T \rightarrow \infty} \mathbf{E} U_T^2, \tag{3.48}$$

where the last equality follows from Lemma 3.4. Taking into account that by (3.2),

$$\mathbf{E} [U_T V_T] = \mathbf{E} [B_{H,\lambda}(T) U_T] - b \mathbf{E} U_T^2,$$

we complete the proof. □

Proposition 3.9. *As $T \rightarrow \infty$,*

$$\begin{pmatrix} Z_T \\ U_T \\ V_T \end{pmatrix} \xrightarrow{d} \mathcal{N}(\mathbf{0}, \Sigma),$$

where

$$\Sigma = \begin{pmatrix} b^{-2} \beta_{H,\lambda,b}^2 & b^{-2} \beta_{H,\lambda,b}^2 & 0 \\ b^{-2} \beta_{H,\lambda,b}^2 & b^{-2} (\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2) & 0 \\ 0 & 0 & \beta_{H,\lambda,b}^2 \end{pmatrix}.$$

In particular,

- Z_T and V_T are asymptotically independent;
- U_T and V_T are asymptotically independent.

Proof. Lemmas 3.3–3.8 together give us the value of the asymptotic covariance matrix. Taking into account the normality of the random vector $(Z_T, V_T, U_T)^\top$, we obtain the desired convergence. \square

3.3. Proof of Theorem 2.1 and convergence (2.12)

Let $(\xi_1, \xi_2, \xi_3) \simeq \mathcal{N}(\mathbf{0}, \Sigma)$, where the matrix Σ is defined in Proposition 3.9.

(i) By Lemma 3.2 and Proposition 3.9,

$$T(\hat{a}_T - a) = b\sigma U_T - \sigma V_T P_T + Q_T \xrightarrow{d} b\sigma \xi_2 - \sigma \xi_3, \quad \text{as } T \rightarrow \infty,$$

where $\xi_2 \simeq \mathcal{N}(0, b^{-2}(\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2))$ and $\xi_3 \simeq \mathcal{N}(0, \beta_{H,\lambda,b}^2)$ are uncorrelated (hence, independent) Gaussian random variables. Therefore, the limiting distribution $b\sigma \xi_2 - \sigma \xi_3$ is zero-mean Gaussian with variance

$$b^2 \sigma^2 b^{-2} (\alpha_{H,\lambda}^2 - \beta_{H,\lambda,b}^2) + \sigma^2 \beta_{H,\lambda,b}^2 = \sigma^2 \alpha_{H,\lambda}^2.$$

Thus, (2.11) is proved.

(ii) According to Lemma 3.1, one has the following representation

$$e^{bt} (\hat{b}_T - b) = \frac{(y_0 + \frac{a}{b} + b\sigma Z_T)^2}{D_T} \frac{\sigma V_T}{y_0 + \frac{a}{b} + b\sigma Z_T} + R_T,$$

where $R_T \rightarrow 0$ a.s. when $T \rightarrow \infty$. By (3.8) and (3.10),

$$\frac{(y_0 + \frac{a}{b} + b\sigma Z_T)^2}{D_T} \rightarrow \frac{\zeta^2}{\frac{1}{2b}\zeta^2} = 2b \quad \text{a.s., as } T \rightarrow \infty.$$

Therefore, we derive from Proposition 3.9 and the Slutsky theorem that

$$e^{bt} (\hat{b}_T - b) \xrightarrow{d} \frac{2b\sigma \xi_3}{y_0 + \frac{a}{b} + b\sigma \xi_1} = \frac{\eta_1}{\eta_2},$$

where $\eta_1 := 2b\sigma \xi_3 \simeq \mathcal{N}(0, 4b^2 \sigma^2 \beta_{H,\lambda,b}^2)$ and $\eta_2 := y_0 + \frac{a}{b} + b\sigma \xi_1 \simeq \mathcal{N}(y_0 + \frac{a}{b}, \sigma^2 \beta_{H,\lambda,b}^2)$ are uncorrelated, hence, independent. \square

A. Asymptotic behavior of drift parameters estimator for the Vasicek model driven by Gaussian process

In this appendix we formulate the main result of Es-Sebaiy and Es-Sebaiy (2021) concerning the asymptotic distribution of the parameter estimators in the Gaussian Vasicek-type model and give the comments which conditions of this paper are satisfied and which are not, therefore it was necessary to modify the respective proofs. So, let $G := \{G_t, t \geq 0\}$ be a centered Gaussian process satisfying the following assumption

(\mathcal{A}_1) There exist constants $c > 0$ and $\gamma \in (0, 1)$ such that for every $s, t \geq 0$,

$$G_0 = 0, \quad \mathbf{E} \left[(G_t - G_s)^2 \right] \leq c|t - s|^{2\gamma}.$$

The Gaussian Vasicek-type process $X = \{X_t, t \geq 0\}$ is defined as the unique (pathwise) solution to

$$X_0 = 0, \quad dX_t = (a + bX_t) dt + dG_t, \quad t \geq 0.$$

where $a \in \mathbb{R}$ and $b > 0$ are considered as unknown parameters. The corresponding least-squares estimators have the form

$$\tilde{b}_T = \frac{\frac{1}{2}T X_T^2 - X_T \int_0^T X_s ds}{T \int_0^T X_s^2 ds - \left(\int_0^T X_s ds\right)^2} \quad \text{and} \quad \tilde{a}_T = \frac{X_T \int_0^T X_s^2 ds - \frac{1}{2}X_T^2 \int_0^T X_s ds}{T \int_0^T X_s^2 ds - \left(\int_0^T X_s ds\right)^2}.$$

The following additional assumptions are required:

(\mathcal{A}_2) There exist $\lambda_G > 0$ and $\eta \in (0, 1)$ such that, as $T \rightarrow \infty$

$$\frac{\mathbf{E}(G_T^2)}{T^{2\eta}} \rightarrow \lambda_G^2. \tag{A.1}$$

(\mathcal{A}_3) There exists a constant $\sigma_G > 0$ such that

$$\lim_{T \rightarrow \infty} \mathbf{E} \left[\left(e^{-bT} \int_0^T e^{bs} dG_s \right)^2 \right] = \sigma_G^2.$$

(\mathcal{A}_4) For all fixed $s \geq 0$,

$$\lim_{T \rightarrow \infty} \mathbf{E} \left(G_s e^{-bT} \int_0^T e^{br} dG_r \right) = 0.$$

(\mathcal{A}_5) For all fixed $s \geq 0$,

$$\lim_{T \rightarrow \infty} \frac{\mathbf{E}(G_s G_T)}{T^\eta} = 0, \quad \lim_{T \rightarrow \infty} \mathbf{E} \left(\frac{G_T}{T^\eta} e^{-bT} \int_0^T e^{br} dG_r \right) = 0.$$

Theorem A.1 (Es-Sebaiy and Es-Sebaiy (2021, Theorem 3.2)). *Assume that (\mathcal{A}_1)–(\mathcal{A}_4) hold. Suppose that $N_1 \sim \mathcal{N}(0, 1)$, $N_2 \sim \mathcal{N}(0, 1)$ and G are independent. Then as $T \rightarrow \infty$,*

$$e^{bT} (\tilde{b}_T - b) \xrightarrow{d} \frac{2b\sigma_G N_2}{\frac{a}{b} + \zeta_\infty},$$

$$T^{1-\eta} (\tilde{a}_T - a) \xrightarrow{d} \lambda_G N_1,$$

where $\zeta_\infty = b \int_0^\infty e^{-bs} G_s ds$. Moreover, if (\mathcal{A}_5) holds, then as $T \rightarrow \infty$,

$$\left(e^{bT} (\tilde{b}_T - b), T^{1-\eta} (\tilde{a}_T - a) \right) \xrightarrow{d} \left(\frac{2b\sigma_G N_2}{\frac{a}{b} + \zeta_\infty}, \lambda_G N_1 \right).$$

Let us analyze whether the conditions (\mathcal{A}_1)–(\mathcal{A}_5) are satisfied for our tempered fractional Vasicek model (2.5). We start with the basic condition (\mathcal{A}_1). The behavior of the variogram function $\mathbf{E}(B_{H,\lambda}(t) - B_{H,\lambda}(s))^2$ of TFBM was recently studied in Mishura and Ralchenko (2024), where the following upper bounds have been established (see Mishura and Ralchenko 2024, Lemma 2):

(i) If $H \in (0, 1)$, then for all $t, s \in \mathbb{R}_+$

$$\mathbf{E} |B_{H,\lambda}(t) - B_{H,\lambda}(s)|^2 \leq C \left(|t - s|^{2H} \wedge 1 \right).$$

(ii) If $H = 1$, then for all $t, s \in \mathbb{R}_+$

$$\mathbf{E} |B_{H,\lambda}(t) - B_{H,\lambda}(s)|^2 \leq C \left(|t - s|^2 |\log |t - s|| \wedge 1 \right).$$

(iii) If $H > 1$, then for all $t, s \in \mathbb{R}_+$

$$\mathbf{E} |B_{H,\lambda}(t) - B_{H,\lambda}(s)|^2 \leq C(|t - s|^2 \wedge 1).$$

Comparison of these bounds with the assumption (\mathcal{A}_1) shows that this assumption holds only for $H \in (0, 1)$. For $H > 1$ one should choose $\gamma = 1$ in this assumption, which is impossible. More careful analysis of the proofs in [Es-Sebaïy and Es-Sebaïy \(2021\)](#) shows that the condition $\gamma < 1$ is substantial for ([Es-Sebaïy and Es-Sebaïy 2021](#), Lemma 2.2), which provides the almost sure convergences

$$\frac{G_T}{T^\delta} \rightarrow 0, \quad \frac{e^{-bt}}{T} \int_0^T |G_t X_t| dt \rightarrow 0, \quad T \rightarrow \infty, \quad (\text{A.2})$$

for any $\gamma < \delta \leq 1$. Based on this convergence, [Es-Sebaïy and Es-Sebaïy \(2021\)](#) derive the convergences of the form (3.4)–(3.7), on which the subsequent study of the estimators is based. Thus, we cannot apply the results of [Es-Sebaïy and Es-Sebaïy \(2021\)](#) directly to the case $G = B_{H,\lambda}$ when $H \geq 1$. However, the almost sure upper bound (2.4) allows us to obtain the first convergence in (A.2) for any $\delta > 0$. This bound also makes it possible to derive the second convergence (see the proof of ([Mishura and Ralchenko 2024](#), Lemma 7)) as well as the convergences (3.4)–(3.7) ([Mishura and Ralchenko 2024](#), Lemma 6), which in turn lead to the strong consistency of the estimators ([Mishura and Ralchenko 2024](#), Theorem 6).

Furthermore, for the case $G = B_{H,\lambda}$, the condition (\mathcal{A}_2) is also violated (for any $H > 0$). Indeed, in view of (2.3), the convergence (A.1) in (\mathcal{A}_2) holds with $\eta = 0$ instead of $\eta \in (0, 1)$. This affects on the behaviour of the estimator \hat{a}_T , which has the following representation:

$$T^{1-\eta}(\hat{a}_T - a) = -\frac{1}{T^\eta} e^{bT} (\tilde{b}_T - b) e^{-bT} \int_0^T X_t dt + \frac{G_T}{T^\eta}.$$

If $\eta > 0$, then the first term of this representation vanishes as $T \rightarrow \infty$, and the second one converges to a normal distribution $\mathcal{N}(0, \lambda_G^2)$. If $\eta = 0$, then both terms have non-trivial limits (in fact, they both converge to normal distributions); hence, the study of the asymptotic behavior of their sum becomes more involved.

The conditions (\mathcal{A}_3) and (\mathcal{A}_4) are satisfied when G is a TFBM. Namely, (\mathcal{A}_3) is verified in Lemma 3.5, and (\mathcal{A}_4) can be checked in a similar way.

Finally, the condition (\mathcal{A}_5) does not hold in the case of TFBM (for all $H > 0$). In particular, for any $s > 0$,

$$\lim_{T \rightarrow \infty} \mathbf{E}[B_{H,\lambda}(s)B_{H,\lambda}(T)] = \frac{1}{2} C_s^2 s^{2H} \neq 0,$$

and, moreover,

$$\begin{aligned} \lim_{T \rightarrow \infty} \mathbf{E} \left[B_{H,\lambda}(T) e^{-bT} \int_0^T e^{br} dB_{H,\lambda}(r) \right] &= \lim_{T \rightarrow \infty} \mathbf{E}[B_{H,\lambda}(T)V_T] \\ &= \lim_{T \rightarrow \infty} \left(\mathbf{E}B_{H,\lambda}^2(T) - b\mathbf{E}[B_{H,\lambda}(T)U_T] \right) = \beta_{H,\lambda,b}^2 \neq 0. \end{aligned}$$

where the limit is computed by (2.3) and (3.48). Thus, both equalities in (\mathcal{A}_5) are not valid. Consequently, the asymptotic independence of the estimators is not guaranteed in the case of TFBM. In Remark 2.2 we explain the correlation between estimators in more detail.

Additionally, compared to [Es-Sebaïy and Es-Sebaïy \(2021\)](#), we do not restrict ourselves with zero initial condition allowing it to be any non-random constant $Y_0 = y_0 \in \mathbb{R}$.

B. Special functions K_ν and ${}_2F_1$

In this appendix, we present the definitions of the function K_ν , which appears in the representation of the covariance function of TFBM (see (2.1)–(2.2)), and the function ${}_2F_1$ from the representation (2.15) for the constant $\beta_{H,\lambda,b}^2$. For further information on this topic, we refer to the book Andrews, Askey, and Roy (1999).

The *modified Bessel function of the second kind* $K_\nu(x)$ has the integral representation

$$K_\nu(x) = \int_0^\infty e^{-x \cosh t} \cosh \nu t \, dt,$$

where $\nu > 0$, $x > 0$. The function $K_\nu(x)$ also has the series representation

$$K_\nu(x) = \frac{\pi}{2} \frac{I_{-\nu}(x) - I_\nu(x)}{\sin(\pi\nu)},$$

where $I_\nu(x) = (\frac{1}{2}|x|)^\nu \sum_{n=0}^\infty \frac{(\frac{1}{2}x)^{2n}}{n! \Gamma(n+1+\nu)}$ is called the *modified Bessel function of the first kind*.

The *Gauss hypergeometric function* ${}_2F_1(a, b; c; x)$ can be defined for complex a , b , c and x . Here, we restrict ourselves to the case of real arguments. Moreover, we assume that $c > b > 0$. In this case, we may define ${}_2F_1(a, b; c; x)$ for $x < 1$ by the following Euler's integral representation (Andrews *et al.* 1999, Theorem 2.2.1):

$${}_2F_1(a, b; c; x) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-xt)^{-a} \, dt.$$

Acknowledgments

The first author is supported by the Swedish Foundation for Strategic Research, grant no. UKR22-0017.

The second author is supported by the Research Council of Finland, decision number 359815.

The first and the second authors acknowledge that the present research is carried out within the frame and support of the ToppForsk project no. 274410 of the Research Council of Norway with the title STORM: Stochastics for Time-Space Risk Models.

References

- Andrews GE, Askey R, Roy R (1999). *Special Functions*, volume 71 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, Cambridge. doi:10.1017/CB09781107325937.
- Azmoodeh E, Mishura Y, Sabzikar F (2022). “How Does Tempering Affect the Local and Global Properties of Fractional Brownian Motion?” *Journal of Theoretical Probability*, **35**(1), 484–527. doi:10.1007/s10959-020-01068-z.
- Belfadli R, Es-Sebajy K, Ouknine Y (2011). “Parameter Estimation for Fractional Ornstein-Uhlenbeck Processes: Non-ergodic Case.” *Frontiers in Science and Engineering*, **1**, 1–16.
- Bishwal JPN (2008). *Parameter Estimation in Stochastic Differential Equations*, volume 1923 of *Lecture Notes in Mathematics*. Springer, Berlin. doi:10.1007/978-3-540-74448-1.
- Chen Y, Hu Y, Wang Z (2017). “Parameter Estimation of Complex Fractional Ornstein-Uhlenbeck Processes with Fractional Noise.” *ALEA. Latin American Journal of Probability and Mathematical Statistics*, **14**(1), 613–629.

- Chen Y, Zhou H (2021). “Parameter Estimation for an Ornstein-Uhlenbeck Process Driven by a General Gaussian Noise.” *Acta Mathematica Scientia. Series B. English Edition*, **41**(2), 573–595. doi:10.1007/s10473-021-0218-x.
- El Machkouri M, Es-Sebaïy K, Ouknine Y (2016). “Least Squares Estimator for Non-ergodic Ornstein-Uhlenbeck Processes Driven by Gaussian Processes.” *Journal of the Korean Statistical Society*, **45**(3), 329–341. doi:10.1016/j.jkss.2015.12.001.
- Es-Sebaïy K, Al-Foraih M, Alazemi F (2021). “Statistical Inference for Nonergodic Weighted Fractional Vasicek Models.” *Modern Stochastics. Theory and Applications*, **8**(3), 291–307. doi:10.15559/21-vmsta176.
- Es-Sebaïy K, EsSebaïy M (2021). “Estimating Drift Parameters in a Non-ergodic Gaussian Vasicek-type Model.” *Statistical Methods & Applications. Journal of the Italian Statistical Society*, **30**(2), 409–436. doi:10.1007/s10260-020-00528-4.
- Gradshteyn IS, Ryzhik IM (2007). *Table of Integrals, Series, and Products*. 7th edition. Academic Press.
- Hu Y, Nualart D (2010). “Parameter Estimation for Fractional Ornstein-Uhlenbeck Processes.” *Statistics & Probability Letters*, **80**(11-12), 1030–1038. doi:10.1016/j.spl.2010.02.018.
- Hu Y, Nualart D, Zhou H (2019). “Parameter Estimation for Fractional Ornstein-Uhlenbeck Processes of General Hurst Parameter.” *Statistical Inference for Stochastic Processes*, **22**(1), 111–142.
- Hu Y, Xi Y (2022). “Parameter Estimation for Threshold Ornstein-Uhlenbeck Processes from Discrete Observations.” *Journal of Computational and Applied Mathematics*, **411**, Paper No. 114264, 17. doi:10.1016/j.cam.2022.114264.
- Iacus SM (2008). *Simulation and Inference for Stochastic Differential Equations. With R Examples*. Springer Series in Statistics. Springer, New York. doi:10.1007/978-0-387-75839-8.
- Jiang H, Dong X (2015). “Parameter Estimation for the Non-stationary Ornstein-Uhlenbeck Process with Linear Drift.” *Statistical Papers*, **56**(1), 257–268. doi:10.1007/s00362-014-0580-z.
- Jiang H, Liu H, Zhou Y (2020). “Asymptotic Properties for the Parameter Estimation in Ornstein-Uhlenbeck Process with Discrete Observations.” *Electronic Journal of Statistics*, **14**(2), 3192–3229. doi:10.1214/20-EJS1738.
- Kawai R (2013). “Local Asymptotic Normality Property for Ornstein-Uhlenbeck Processes with Jumps under Discrete Sampling.” *Journal of Theoretical Probability*, **26**(4), 932–967. doi:10.1007/s10959-012-0455-y.
- Kleptsyna ML, Le Breton A (2002). “Statistical Analysis of the Fractional Ornstein-Uhlenbeck Type Process.” *Statistical Inference for Stochastic Processes. An International Journal Devoted to Time Series Analysis and the Statistics of Continuous Time Processes and Dynamical Systems*, **5**(3), 229–248. doi:10.1023/A:1021220818545.
- Kubilius K, Mishura Y, Ralchenko K (2017). *Parameter Estimation in Fractional Diffusion Models*, volume 8 of *Bocconi & Springer Series*. Bocconi University Press, Milan; Springer, Cham. doi:10.1007/978-3-319-71030-3.
- Kubilius K, Mishura Y, Ralchenko K, Seleznev O (2015). “Consistency of the Drift Parameter Estimator for the Discretized Fractional Ornstein-Uhlenbeck Process with Hurst Index $H \in (0, \frac{1}{2})$.” *Electronic Journal of Statistics*, **9**(2), 1799–1825.

- Kukush A, Mishura Y, Ralchenko K (2017). “Hypothesis Testing of the Drift Parameter Sign for Fractional Ornstein-Uhlenbeck Process.” *Electronic Journal of Statistics*, **11**(1), 385–400. doi:10.1214/17-EJS1237.
- Kutoyants YA (2004). *Statistical Inference for Ergodic Diffusion Processes*. Springer Series in Statistics. Springer-Verlag London, Ltd., London. doi:10.1007/978-1-4471-3866-2.
- Li S, Dong Y (2018). “Parametric Estimation in the Vasicek-Type Model Driven by Sub-Fractional Brownian Motion.” *Algorithms*, **11**, 197. doi:10.3390/a11120197.
- Liptser RS, Shiryaev AN (2001). *Statistics of Random Processes. II. Applications*, volume 6 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin.
- Lohvinenko S, Ralchenko K (2017). “Maximum Likelihood Estimation in the Fractional Vasicek Model.” *Lithuanian Journal of Statistics*, **56**(1), 77–87.
- Lohvinenko S, Ralchenko K (2018). “Asymptotic Distribution of Maximum Likelihood Estimator in Fractional Vasicek Model.” *Theory of Probability and Mathematical Statistics*, **99**, 134–151.
- Lohvinenko S, Ralchenko K (2019). “Maximum Likelihood Estimation in the Non-ergodic Fractional Vasicek Model.” *Modern Stochastics. Theory and Applications*, **6**(3), 377–395. doi:10.15559/19-vmsta140.
- Lohvinenko S, Ralchenko K, Zhuchenko O (2016). “Asymptotic Properties of Parameter Estimators in Fractional Vasicek Model.” *Lithuanian Journal of Statistics*, **55**(1), 102–111.
- Meerschaert MM, Sabzikar F (2013). “Tempered Fractional Brownian Motion.” *Statistics & Probability Letters*, **83**(10), 2269–2275. doi:10.1016/j.spl.2013.06.016.
- Meerschaert MM, Sabzikar F (2014). “Stochastic Integration for Tempered Fractional Brownian Motion.” *Stochastic Processes and Their Applications*, **124**(7), 2363–2387. doi:10.1016/j.spa.2014.03.002.
- Mendy I (2013). “Parametric Estimation for Sub-fractional Ornstein-Uhlenbeck Process.” *Journal of Statistical Planning and Inference*, **143**(4), 663–674. doi:10.1016/j.jspi.2012.10.013.
- Mishura Y, Ralchenko K (2017). “Drift Parameter Estimation in the Models Involving Fractional Brownian Motion.” In *Modern Problems of Stochastic Analysis and Statistics*, volume 208 of *Springer Proceedings in Mathematics & Statistics*, pp. 237–268. Springer, Cham.
- Mishura Y, Ralchenko K (2024). “Asymptotic Growth of Sample Paths of Tempered Fractional Brownian Motions, with Statistical Applications to Vasicek-type Models.” *Fractal and Fractional*, **8**(2:79).
- Mishura Y, Ralchenko K, Shklyar S (2023). “Gaussian Volterra Processes: Asymptotic Growth and Statistical Estimation.” *Theory of Probability and Mathematical Statistics*, (108), 149–167. doi:10.1090/tpms/1190.
- Moers M (2012). “Hypothesis Testing in a Fractional Ornstein-Uhlenbeck Model.” *International Journal of Stochastic Analysis*, pp. Art. ID 268568, 23. doi:10.1155/2012/268568.
- Nourdin I, Tran TTD (2019). “Statistical Inference for Vasicek-type Model Driven by Hermite Processes.” *Stochastic Processes and Their Applications*, **129**(10), 3774–3791. doi:10.1016/j.spa.2018.10.005.
- Prykhodko O, Ralchenko K (2024). “Asymptotic Normality of Estimators for All Parameters in the Vasicek Model by Discrete Observations.” *Theory of Probability and Mathematical Statistics*, **111**.

- Shen G, Yin X, Yan L (2016). “Least Squares Estimation for Ornstein-Uhlenbeck Processes Driven by the Weighted Fractional Brownian Motion.” *Acta Mathematica Scientia. Series B. English Edition*, **36**(2), 394–408. doi:10.1016/S0252-9602(16)30008-X.
- Shen G, Yu Q (2019). “Least Squares Estimator for Ornstein-Uhlenbeck Processes Driven by Fractional Lévy Processes from Discrete Observations.” *Statistical Papers*, **60**(6), 2253–2271. doi:10.1007/s00362-017-0918-4.
- Shimizu Y (2012). “Local Asymptotic Mixed Normality for Discretely Observed Non-recurrent Ornstein-Uhlenbeck Processes.” *Annals of the Institute of Statistical Mathematics*, **64**(1), 193–211. doi:10.1007/s10463-010-0307-4.
- Tanaka K (2013). “Distributions of the Maximum Likelihood and Minimum Contrast Estimators Associated with the Fractional Ornstein-Uhlenbeck Process.” *Statistical Inference for Stochastic Processes. An International Journal Devoted to Time Series Analysis and the Statistics of Continuous Time Processes and Dynamical Systems*, **16**(3), 173–192. doi:10.1007/s11203-013-9085-y.
- Tanaka K (2015). “Maximum Likelihood Estimation for the Non-ergodic Fractional Ornstein-Uhlenbeck Process.” *Statistical Inference for Stochastic Processes. An International Journal Devoted to Time Series Analysis and the Statistics of Continuous Time Processes and Dynamical Systems*, **18**(3), 315–332. doi:10.1007/s11203-014-9110-9.
- Tanaka K, Xiao W, Yu J (2020). “Maximum Likelihood Estimation for the Fractional Vasicek Model.” *Econometrics*, **8**(3), 32.
- Tang CY, Chen SX (2009). “Parameter Estimation and Bias Correction for Diffusion Processes.” *Journal of Econometrics*, **149**(1), 65–81. doi:10.1016/j.jeconom.2008.11.001.
- Tudor CA, Viens FG (2007). “Statistical Aspects of the Fractional Stochastic Calculus.” *The Annals of Statistics*, **35**(3), 1183–1212.
- Wang J, Xiao X, Li C (2023). “Least Squares Estimations for Approximate Fractional Vasicek Model Driven by a Semimartingale.” *Mathematics and Computers in Simulation*, **208**, 207–218. doi:10.1016/j.matcom.2023.01.015.
- Xiao W, Yu J (2019a). “Asymptotic Theory for Estimating Drift Parameters in the Fractional Vasicek Model.” *Econometric Theory*, **35**(1), 198–231. doi:10.1017/S0266466618000051.
- Xiao W, Yu J (2019b). “Asymptotic Theory for Rough Fractional Vasicek Models.” *Economics Letters*, **177**, 26–29. doi:10.1016/j.econlet.2019.01.020.
- Zang Q, Zhang L (2019). “Asymptotic Behaviour of the Trajectory Fitting Estimator for Reflected Ornstein-Uhlenbeck Processes.” *Journal of Theoretical Probability*, **32**(1), 183–201. doi:10.1007/s10959-017-0796-7.

Affiliation:

Yuliya Mishura

Department of Probability, Statistics and Actuarial Mathematics

Taras Shevchenko National University of Kyiv

64/13, Volodymyrs'ka St., 01601 Kyiv, Ukraine

E-mail: yuliyamishura@knu.ua

Division of Mathematics and Physics

Mälardalen University

721 23 Västerås, Sweden

E-mail: yuliia.mishura@mdu.se

Kostiantyn Ralchenko
Department of Probability, Statistics and Actuarial Mathematics
Taras Shevchenko National University of Kyiv
64/13, Volodymyrs'ka St., 01601 Kyiv, Ukraine
E-mail: kostiantynralchenko@knu.ua

School of Technology and Innovations
University of Vaasa
P.O. Box 700, Wolffintie 34, FIN-65101 Vaasa, Finland
E-mail: kostiantyn.ralchenko@uwasa.fi

Olena Dehtiar
Department of Probability, Statistics and Actuarial Mathematics
Taras Shevchenko National University of Kyiv
64/13, Volodymyrs'ka St., 01601 Kyiv, Ukraine
E-mail: dehtiar.olena@knu.ua